



MOK (Phospho-Tyr161) Monoclonal Antibody

Catalog No	BYmab-10345
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	MOK RAGE RAGE1
Protein Name	MOK (Phospho-Tyr161)
Immunogen	Synthesized peptide derived from human MOK (Phospho-Tyr161)
Specificity	This antibody detects endogenous phospho levels of MOK (Phospho-Tyr161) at Human:Y161, Mouse:Y161, Rat:Y161
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,lgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year MAPK/MAK/MRK overlapping kinase (EC 2.7.11.22;MOK protein kinase;Renal tumor antigen 1;RAGE-1)
	MAPK/MAK/MRK overlapping kinase (EC 2.7.11.22;MOK protein kinase;Renal
Synonyms	MAPK/MAK/MRK overlapping kinase (EC 2.7.11.22;MOK protein kinase;Renal tumor antigen 1;RAGE-1)
Synonyms Observed Band	MAPK/MAK/MRK overlapping kinase (EC 2.7.11.22;MOK protein kinase;Renal tumor antigen 1;RAGE-1) 48kD
Synonyms Observed Band Cell Pathway	MAPK/MAK/MRK overlapping kinase (EC 2.7.11.22;MOK protein kinase;Renal tumor antigen 1;RAGE-1) 48kD Cytoplasm . Cell projection, cilium . Nucleus . Expressed in heart, brain, lung, kidney, and pancreas, and at very low levels in placenta, liver and skeletal muscle. Detected in retina. catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Phosphorylation appears to increase the enzymatic activity.,function:Able to phosphorylate several exogenous substrates and to
Synonyms Observed Band Cell Pathway Tissue Specificity	MAPK/MAK/MRK overlapping kinase (EC 2.7.11.22;MOK protein kinase;Renal tumor antigen 1;RAGE-1) 48kD Cytoplasm . Cell projection, cilium . Nucleus . Expressed in heart, brain, lung, kidney, and pancreas, and at very low levels in placenta, liver and skeletal muscle. Detected in retina. catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Phosphorylation appears to increase the enzymatic activity.,function:Able to phosphorylate several exogenous substrates and to undergo autophosphorylation.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. CMGC Ser/Thr protein kinase family. CDC2/CDKX subfamily.,similarity:Contains 1 protein kinase domain.,tissue specificity:Expressed in heart, brain, lung, kidney, and pancreas, and at very low

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询




transcription factor 2 (Cdx2) protein. The encoded protein, which is localized to epithelial cells in the intestinal crypt, may play a role in growth arrest and differentiation of cells of upper crypt and lower villus regions. Multiple alternatively spliced transcript variants encoding different isoforms have been observed for this gene. [provided by RefSeq, Dec 2012],
Avoid repeated freezing and thawing!

matters needing attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using MOK (Phospho-Tyr161) Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658