



NEDD4L (phospho-Ser342) mouse mAb

Catalog No	BYmab-10387
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	NEDD4L KIAA0439 NEDL3
Protein Name	NEDD4L (Ser342)
Immunogen	Synthesized phosho peptide around human NEDD4L (Ser342)
Specificity	This antibody detects endogenous levels of Human Mouse Rat NEDD4L (phospho-Ser342)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	E3 ubiquitin-protein ligase NEDD4-like (EC 6.3.2.-) (NEDD4.2) (Nedd4-2)
Observed Band	110-120kD
Cell Pathway	Cytoplasm . Golgi apparatus . Endosome, multivesicular body . May be recruited to exosomes by NDFIP1.
Tissue Specificity	Ubiquitously expressed, with highest levels in prostate, pancreas, and kidney (PubMed:14615060, PubMed:15496141, PubMed:19664597). Expressed in melanocytes (PubMed:23999003).
Function	function:E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Inhibits TGF-beta signaling by triggering SMAD2 and TGF-beta1 ubiquitination and proteasome-dependent degradation. Promotes ubiquitination and internalization of various plasma membrane channels such as ENaC, Nav1.2, Nav1.3, Nav1.5, Nav1.7, Nav1.8, Kv1.3, EAAT1 or CLC5. Promotes ubiquitination and degradation of SGK.,induction:By androgens in prostate, and by albumin in kidney.,pathway:Protein modification; protein ubiquitination.,PTM:Phosphorylated by SGK or PKA; which impairs interaction with SCNN. Interaction with YWHAH inhibits

Nanjing BYabscience technology Co.,Ltd



dephosphorylation.,similarity:Contains 1 C2 domain.,similarity:Contains 1 HECT (E6AP-type E3 ubiquitin-protein ligase) domain.,similarity:Contains 4 WW domains.,subunit:Int

Background

This gene encodes a member of the Nedd4 family of HECT domain E3 ubiquitin ligases. HECT domain E3 ubiquitin ligases transfer ubiquitin from E2 ubiquitin-conjugating enzymes to protein substrates, thus targeting specific proteins for lysosomal degradation. The encoded protein mediates the ubiquitination of multiple target substrates and plays a critical role in epithelial sodium transport by regulating the cell surface expression of the epithelial sodium channel, ENaC. Single nucleotide polymorphisms in this gene may be associated with essential hypertension. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Mar 2012],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using NEDD4L (phospho-Ser342) mouse mAb



Nanjing BYabs science technology Co.,Ltd