




TAZ (phospho-Ser89) mouse mAb

Catalog No	BYmab-10414
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	WWTR1 TAZ
Protein Name	TAZ (Ser89)
Immunogen	Synthesized phosho peptide around human TAZ (Ser89)
Specificity	This antibody detects endogenous levels of Human Mouse Rat TAZ (phospho-Ser89)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
D i to .	≥90%
Purity	290%
Storage Stability	-20°C/1 year
Storage Stability	-20°C/1 year WW domain-containing transcription regulator protein 1 (Transcriptional
Storage Stability Synonyms	-20°C/1 year WW domain-containing transcription regulator protein 1 (Transcriptional coactivator with PDZ-binding motif) 44kD Nucleus . Cytoplasm . Cell membrane . Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies (By similarity). When phosphorylated, is retained in the cytoplasm by YWHAZ (By similarity). Can be retained in the nucleus by MED15 (PubMed:18568018). Localized in the cytoplasm in areas of epithelial cell high density (PubMed:21145499). At blastocyst stage expressed in the nucleus in
Storage Stability Synonyms Observed Band	-20°C/1 year WW domain-containing transcription regulator protein 1 (Transcriptional coactivator with PDZ-binding motif) 44kD Nucleus . Cytoplasm . Cell membrane . Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies (By similarity). When phosphorylated, is retained in the cytoplasm by YWHAZ (By similarity). Can be retained in the nucleus by MED15 (PubMed:18568018). Localized in the cytoplasm in areas of epithelial cell high density (PubMed:21145499). At blastocyst stage expressed in the nucleus in trophectodermal cells, however expressed in the cytoplasm in the inner cell mass
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year WW domain-containing transcription regulator protein 1 (Transcriptional coactivator with PDZ-binding motif) 44kD Nucleus . Cytoplasm . Cell membrane . Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies (By similarity). When phosphorylated, is retained in the cytoplasm by YWHAZ (By similarity). Can be retained in the nucleus by MED15 (PubMed:18568018). Localized in the cytoplasm in areas of epithelial cell high density (PubMed:21145499). At blastocyst stage expressed in the nucleus in trophectodermal cells, however expressed in the cytoplasm in the inner cell mass (By similarity) Highly expressed in kidney, heart, placenta and lung. Expressed in the thyroid

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







coactivator.,PTM:Phosphorylated. Phosphorylation results in the inhibition of transcriptional coactivation through YWHAZ-mediated nuclear export.,similarity:Contains 1 WW domain.,subcellular location:Concentrates along specific portions of the plasma membrane, and accumulates in punctate nuclear bodies. When phosphorylated, is retained in cytoplasm by YWHAZ.,subunit:Binds to SLC9A3R2 via the PDZ motif at the plasma membrane. Binds to YWHAZ in vivo and in vitro through the phosphoserine-binding motif RSHSSP.,tissue specificity:Highly expressed in kidney, heart, placenta and lung.,

Background


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TAZ (phospho-Ser89) mouse mAb

Nanjing BYabscience technology Co.,Ltd