



TIFA(Phospho-Thr9) mouse mAb

| Catalog No BYmab-10483 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name TIFA T2BP Protein Name TIFA(Phospho-Thr9) Immunogen Synthesized peptide derived from human TIFA(Phospho-Thr9) Specificity This antibody detects endogenous levels of TIFA(Phospho-Thr9) at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm Colocalizes with Iysosomal marker LAMP2 following homooligomerization and subsequent activation. Tissue Specificity Function Background | | |
|---|---------------------------|---|
| Reactivity Human; Mouse;Rat Applications WB Gene Name TIFA T2BP Protein Name ITIFA(Phospho-Thr9) Immunogen Synthesized peptide derived from human TIFA(Phospho-Thr9) Specificity This antibody detects endogenous levels of TIFA(Phospho-Thr9) at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing! | Catalog No | BYmab-10483 |
| Applications WB Gene Name TIFA T2BP Protein Name TIFA(Phospho-Thr9) Immunogen Synthesized peptide derived from human TIFA(Phospho-Thr9) Specificity This antibody detects endogenous levels of TIFA(Phospho-Thr9) at Human, Mouse, Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation. Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing! | Isotype | IgG |
| Gene Name TIFA T2BP Protein Name TIFA(Phospho-Thr9) Immunogen Synthesized peptide derived from human TIFA(Phospho-Thr9) Specificity This antibody detects endogenous levels of TIFA(Phospho-Thr9) at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing! | Reactivity | Human; Mouse;Rat |
| Protein Name TIFA(Phospho-Thr9) Immunogen Synthesized peptide derived from human TIFA(Phospho-Thr9) Specificity This antibody detects endogenous levels of Mouse, Rat TIFA(Phospho-Thr9) at Human, Mouse, Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Applications | WB |
| Immunogen Synthesized peptide derived from human TIFA(Phospho-Thr9) Specificity This antibody detects endogenous levels of TIFA(Phospho-Thr9) at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein) PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with Iysosomal marker LAMP2 following homooligomerization and subsequent activation | Gene Name | TIFA T2BP |
| Specificity This antibody detects endogenous levels of TIFA(Phospho-Thr9) at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with Iysosomal marker LAMP2 following homooligomerization and subsequent activation. Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing! | Protein Name | TIFA(Phospho-Thr9) |
| Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Matters needing Avoid repeated freezing and thawing! | Immunogen | Synthesized peptide derived from human TIFA(Phospho-Thr9) |
| Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with Iysosomal marker LAMP2 following homooligomerization and subsequent activation. Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing! | Specificity | This antibody detects endogenous levels of TIFA(Phospho-Thr9) at Human, Mouse,Rat |
| Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Matters needing Avoid repeated freezing and thawing! | Formulation | Liquid in PBS containing 50% glycerol, and 0.138% sodium azide. |
| affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Source | Monoclonal, Mouse,IgG |
| Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Purification | · |
| Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Dilution | WB 1:500-2000 |
| Storage Stability -20°C/1 year TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Concentration | 1 mg/ml |
| Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Purity | ≥90% |
| MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Storage Stability | -20°C/1 year |
| Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing! | Synonyms | MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) |
| homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing! | Observed Band | |
| Function Background matters needing Avoid repeated freezing and thawing! | Cell Pathway | Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation |
| Background matters needing Avoid repeated freezing and thawing! | Tissue Specificity | |
| matters needing Avoid repeated freezing and thawing! | Function | |
| matters needing Avoid repeated freezing and thawing! attention | Background | |
| | matters needing attention | Avoid repeated freezing and thawing! |

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TIFA(Phospho-Thr9) mouse mAb

Nanjing BYabscience technology Co.,Ltd