



## Gz-α (Phospho-Ser16) mouse mAb

Catalog No BYmab-10536   Isotype IgG   Reactivity Human; Mouse;Rat   Applications WB   Gene Name GNAZ   Protein Name Gz-a (Phospho-Ser16)   Immunogen Synthesized peptide derived from human Gz-a (Phospho-Ser16)   Specificity This antibody detects endogenous levels of Gz-a (Phospho-Ser16) at Human, Mouse,Rat   Formulation Liquid in PBS containing 50% glycerol, and 0.192% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Gaunien nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)   Observed Band 41kD   Call Pathway Membrane; Lipid-anchor.   Function function:Guanien nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling menulsedhile binding site, and antime, activation in the alpha chain contains the guanine nucleotide binding site, act and antime, act and antinstruction in maintating the orige of the sins binfantly G(t/01/2		
ReactivityHuman; Mouse;RatApplicationsWBGene NameGNAZProtein NameGz- a (Phospho-Ser16)ImmunogenSynthesized peptide derived from human Gz- a (Phospho-Ser16)SpecificityThis antibody detects endogenous levels of Gz- a (Phospho-Ser16) at Human, Mouse, RatFormulationLiquid in PBS containing 50% glycerol, and 0.192% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsGuanne nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Functionfunction:Guanne nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarly Belongs to the G-Alpha family, G(10/V2) subfamily, signal and gamma. The alpha chain contains the guanne mucleotide-binding site.BackgroundG protein subout alpha z(GNA2) Homo sapiensThe protein encoded by this gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systems, This encoded protein may play a role in maintaining the lonic balance or perlymphatic and endolymphatic cochear fluids. provided by RefSeq, Jul 2008 .<	Catalog No	BYmab-10536
ApplicationsWBGene NameGNAZProtein NameG2-α (Phospho-Ser16)ImmunogenSynthesized peptide derived from human Gz-α (Phospho-Ser16)SpecificityThis antibody detects endogenous levels of Gz-α (Phospho-Ser16) at Human, Mouse, RatFormulationLiquid in PBS containing 50% glycerol, and 0.192% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsGuanne nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarity-Belongs to the G-alpha family. Gl( <i>io</i> /Iz) subfamily, subfamily, subfamily, Belong of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site,.BackgroundG protein subunit alpha z (SNAZ) Homo sapiens gene is a member of a G protein sitemal gamma. The alpha chain contains the guanine nucleotide binding site,.BackgroundG protein subunit alpha z (SNAZ) Homo sapiens gene is a member of a G protein sitemal gamma. The alpha chain contains the guanine nucleotide binding site,.	Isotype	lgG
Gene Name GNAZ   Protein Name Gz- α (Phospho-Ser16)   Immunogen Synthesized peptide derived from human Gz- α (Phospho-Ser16)   Specificity This antibody detects endogenous levels of Gz- α (Phospho-Ser16) at Human, Mouse, Rat   Formulation Liquid in PBS containing 50% glycerol, and 0.192% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Guanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)   Observed Band 41kD   Cell Pathway Membrane; Lipid-anchor.   Tissue Specificity Brain,PNS,   Function function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarity.Belongs to the G-alpha family. Gl(/ol/2) subfamily, subunit.G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site,.   Background G protein subunit alpha 2(GNAZ) Horo sapiens The protein encoded by this gene is a member of a G protein subamily finat m	Reactivity	Human; Mouse;Rat
Protein Name Gz-α (Phospho-Ser16)   Immunogen Synthesized peptide derived from human Gz-α (Phospho-Ser16)   Specificity This antibody detects endogenous levels of Gz-α (Phospho-Ser16) at Human, Mouse, Rat   Formulation Liquid in PBS containing 50% glycerol, and 0.192% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Guanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)   Observed Band 41kD   Cell Pathway Membrane; Lipid-anchor.   Tissue Specificity Brain,PNS,   Function function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarity:Belongs to the G-alpha family, G(i/ol/2) subfamily, subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide-binding site.   Background G protein subunit alpha z(GNAZ) Homo sapiens The protein encoded by this gene is a member of a G protelymphatic and eolymphatic cochiear fluids.	Applications	WB
ImmunogenSynthesized peptide derived from human Gz- a (Phospho-Ser16)SpecificityThis antibody detects endogenous levels of Gz- a (Phospho-Ser16) at Human, Mouse,RatFormulationLiquid in PBS containing 50% glycerol, and 0.192% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsGuanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarity:Belongs to the G-alpha family. (Gl/ol/12) subfamily., subunit:G proteins are composed of 3 units, alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding ste.,BackgroundG protein subunit alpha z(GNAZ) Homo sagiens gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systems. This encoded protein may play a role in mantaining the ionic balance of perilymphatic cochlear fluids. Iprovided by RefSeq, Jul 2008],	Gene Name	GNAZ
Specificity This antibody detects endogenous levels of Gz- a (Phospho-Ser16) at Human, Mouse, Rat   Formulation Liquid in PBS containing 50% glycerol, and 0.192% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Guanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)   Observed Band 41kD   Cell Pathway Membrane; Lipid-anchor.   Tissue Specificity Brain,PNS,   Function function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarity:Belongs to the G-alpha family. G(i/o/l/2) subarily, subunit:G proteins are compaced sto the G-alpha family. G(i/o/l/2) subarily, subunit:G proteins are contains the guanine nucleotide binding site, and gamma. The alpha chain contains the guanine nucleotide binding site, and gamma. The alpha chain contains the guanine nucleotide binding site, and gamma. The alpha chain contains the guanine nucleotide binding site, and gamma. The alpha chain contains the guanine nucleotide binding site, and gamma. The alpha chain contains the guanine nucleotide binding site, and gamma. The alpha chain contains the guanine nucleotide binding site, and gamma.	Protein Name	Gz-α (Phospho-Ser16)
Mouse,RatFormulationLiquid in PBS containing 50% glycerol, and 0.192% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsGuanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarity:Belongs to the G-alpha family. G(i/ol/t2) subfamily., subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha zG(NAZ) Homo sapiens maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. provided by RefSeq, Jul 2008],	Immunogen	Synthesized peptide derived from human Gz- $\alpha$ (Phospho-Ser16)
SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsGuanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems, similarity:Belongs to the G-alpha family. G(i/o/l/z) subfamily, subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide subing siteBackgroundG protein subunit alpha z(GNAZ) Homo sapiens maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. provided by RefSeq, Jul 2008],	Specificity	
PurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsGuanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction: Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems., similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily, subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide by Brain, PNS,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens metrices signal transduction in proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site., Dace and gamma. The alpha chain contains the guanine nucleotide of protein subfamily that mediates signal transduction in proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site., Imaintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. provided by RefSeq, Jul 2008],	Formulation	Liquid in PBS containing 50% glycerol, and 0.192% sodium azide.
affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Guanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)   Observed Band 41kD   Cell Pathway Membrane; Lipid-anchor.   Tissue Specificity Brain,PNS,   Function function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/lt/2) subfamily.,subunit:G proteins are composed of 3 units; alpha, befa and gamma. The alpha chain contains the guanine nucleotide binding site.,   Background G protein subunit alpha z(GNAZ) Homo sapiens The protein encoded by this gene is a member of a G protein subfamily that mediates signal transduction in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Guanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)   Observed Band 41kD   Cell Pathway Membrane; Lipid-anchor.   Tissue Specificity Brain,PNS,   Function function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems., similarity:Belongs to the G-alpha family. G(v/l/2) subfamilysubunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,   Background G protein subunit alpha z (GNAZ) Homo sapiens   Background G protein subunit alpha z (GNAZ) Homo sapiens   In encoded by RefSeq, Jul 2008], This encoded protein may play a role in maintaining the ionic balance of perlymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Purification	
Purity≥90%Storage Stability-20°C/1 yearSynonymsGuanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/oft/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systms. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Dilution	WB 1:500-2000
Storage Stability-20°C/1 yearSynonymsGuanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systems. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Concentration	1 mg/ml
SynonymsGuanine nucleotide-binding protein G(z) subunit alpha (G(x) alpha chain) (Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Purity	≥90%
(Gz-alpha)Observed Band41kDCell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Storage Stability	-20°C/1 year
Cell PathwayMembrane; Lipid-anchor.Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems., similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily., subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens menture is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systems. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Synonyms	
Tissue SpecificityBrain,PNS,Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens pertussis toxin-insensitive systems. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Observed Band	41kD
Functionfunction:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systms. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Cell Pathway	Membrane; Lipid-anchor.
modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site.,BackgroundG protein subunit alpha z(GNAZ) Homo sapiens gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systms. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Tissue Specificity	Brain,PNS,
gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systms. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008],	Function	modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain
Nanjing BYabscience technology Co.,Ltd	Background	gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systms. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids.
		Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





## matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

SY-SY5Y 170---92---70---53---42---33---22---14---9--- Western Blot analysis of various cells using Gz-  $\alpha$  (Phospho-Ser16) mouse mAb

Nanjing BYabscience technology Co.,Ltd