



IRE1 Phospho-thr973 mouse mAb

Catalog No BYmab-10569 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name ERN1 IRE1 Protein Name IRE1 thr973 Immunogen Synthesized peptide derived from human IRE1 thr973 Specificity This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (InRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue.		
Reactivity Human; Mouse;Rat Applications WB Gene Name ERN1 IRE1 Protein Name IRE1 thr973 Immunogen Synthesized peptide derived from human IRE1 thr973 Specificity This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein_cofactor.Magnesium_enzyme regulation.The kinase domain is activated by trans-autophosphonylation. Kinase activity is required for activation of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBPI mRNA to generate a new C-terminus. Convertion it into a potent unfolded-orotein response	Catalog No	BYmab-10569
Applications Gene Name ERN1 IRE1 Protein Name IRE1 thr973 Immunogen Synthesized peptide derived from human IRE1 thr973 Specificity This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein_cofactor:Magnesium_enzyme regulation:The kinase domain is activated by trans-autophosphonylation. Kinase activity is required for activation of the endoplasmic retriculum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus. convertion it into a potent unfolded-protein response	Isotype	IgG
Gene Name ERN1 IRE1 Protein Name IRE1 thr973 Immunogen Synthesized peptide derived from human IRE1 thr973 Specificity This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein, cofactor:Magnesium, enzyme regulation: The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain	Reactivity	Human; Mouse;Rat
Protein Name IRE1 thr973 Immunogen Synthesized peptide derived from human IRE1 thr973 Specificity This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein, cofactor:Magnesium, enzyme regulation:The kinase activation of the endoribonuclease domain. function:Senses unfolded proteins in the lumen of the endoribonuclease domain, function:Senses unfolded proteins in the lumen of the endoribonuclease and e	Applications	WB
Immunogen Synthesized peptide derived from human IRE1 thr973 Specificity This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein, cofactor:Magnesium, enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain, function:Senses unfolded protein in the lumen of the endoribonuclease domain, function:Senses unfolded protein restoring auto-activation. The active en	Gene Name	ERN1 IRE1
Specificity This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein, cofactor:Magnesium, enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain, function:Senses unfolded proteins in the lumen of the endoribonuclease domain, function:Senses unfolded proteins in the lumen of the endoribonuclease domain which leads to enzyme auto-activation. The extraorion reticulum via its N-terminal domain which leads to en	Protein Name	IRE1 thr973
Formulation Liquid in PBS containing 50% glycerol, and 0.225% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein., cofactor:Magnesium, enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain spices XBPI mRNA to generate a new C-terminus, converting it into a optent unfolded-protein response	Immunogen	Synthesized peptide derived from human IRE1 thr973
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity: ATP + a protein = ADP + a phosphoprotein, cofactor: Magnesium, enzyme regulation: The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain, function: Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to energiate a new C-terminus. converting it into a potent unfolded-protein response	Specificity	This antibody detects endogenous levels of IRE1 thr973 at Human, Mouse,Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein., cofactor:Magnesium., enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain, function:Senses unfolded proteins in the lumen of the endoribonuclease domain, function:Senses unfolded proteins in the lumen of the endoribonuclease domain. function: Senses unfolded proteins in the lumen of the endoribonuclease domain. function: Senses unfolded proteins in the lumen of the endoribonuclease domain. function: Senses unfolded proteins in the lumen of the endoribonuclease domain. function: Senses unfolded proteins in the lumen of the endoribonuclease domain. function: Senses unfolded proteins in the lumen of the endoribonuclease domain. function: Senses unfolded proteins in the lumen of the endoribonuclease domain. function: Senses unfolded proteins in the lumen of the endoribonuclease domain splices XBP1 mRNA to quentate a new C-terminus. convertino it into a potent unfolded-protein response	Formulation	Liquid in PBS containing 50% glycerol, and 0.225% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain, function: Senses unfolded proteins in the lumen of the endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain, function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus. converting it into a potent unfolded-protein response	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane ; Single-pass type I membrane protein . Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain. function:Senses unfolded proteins in the lumen of the endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Concentration	1 mg/ml
Synonyms Serine/threonine-protein kinase/endoribonuclease IRE1 (Endoplasmic reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Purity	≥90%
reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1); Endoribonuclease (EC 3.1.26)] Observed Band 107kD Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Storage Stability	-20°C/1 year
Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Ubiquitously expressed. High levels observed in pancreatic tissue. Function catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Synonyms	reticulum-to-nucleus signaling 1) (Inositol-requiring protein 1) (hIRE1p) (Ire1-alpha) (IRE1a) [Includes: Serine/threonine-protein kinase (EC 2.7.11.1);
Tissue Specificity Ubiquitously expressed. High levels observed in pancreatic tissue. Catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Observed Band	107kD
catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Cell Pathway	Endoplasmic reticulum membrane ; Single-pass type I membrane protein .
phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response	Tissue Specificity	Ubiquitously expressed. High levels observed in pancreatic tissue.
	Function	phosphoprotein.,cofactor:Magnesium.,enzyme regulation:The kinase domain is activated by trans-autophosphorylation. Kinase activity is required for activation of the endoribonuclease domain.,function:Senses unfolded proteins in the lumen of the endoplasmic reticulum via its N-terminal domain which leads to enzyme auto-activation. The active endoribonuclease domain splices XBP1 mRNA to generate a new C-terminus, converting it into a potent unfolded-protein response

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	apoptosis.,PTM:Autophosphorylated.,similarity:Belongs to the protein kinase superfamily. Ser/Thr protein kinase family.,similarity:Contains 1 KEN domain.,similarity:Contains 1 protein kinase domain.,subunit:Homodimer; disulfide-linked. Dimer formation is driven by hydrophobic interactions within the N-terminal luminal domains
Background	The protein encoded by this gene is the ER to nucleus signalling 1 protein, a human homologue of the yeast Ire1 gene product. This protein possesses intrinsic kinase activity and an endoribonuclease activity and it is important in altering gene expression as a response to endoplasmic reticulum-based stress signals. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658