



# FOG-2 Monoclonal Antibody

<b>Catalog No</b>	BYmab-10576
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	ZFPM2
<b>Protein Name</b>	Zinc finger protein ZFPM2
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human ZFPM2. AA range:921-970
<b>Specificity</b>	FOG-2 Monoclonal Antibody detects endogenous levels of FOG-2 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	ZFPM2; FOG2; ZNF89B; Zinc finger protein ZFPM2; Friend of GATA protein 2; FOG-2; Friend of GATA 2; hFOG-2; Zinc finger protein 89B; Zinc finger protein multitype 2
<b>Observed Band</b>	128kD
<b>Cell Pathway</b>	Nucleus .
<b>Tissue Specificity</b>	Widely expressed at low level.
<b>Function</b>	disease:Defects in ZFPM2 are the cause of diaphragmatic hernia 3 (DIH3) [MIM:610187]; a form of congenital diaphragmatic hernia (CDH). CDH refers to a group of congenital defects in the structural integrity of the diaphragm associated with often lethal pulmonary hypoplasia and pulmonary hypertension.;disease:Defects in ZFPM2 may be a cause of tetralogy of Fallot (TOF) [MIM:187500]. TOF is a congenital heart anomaly which consists of pulmonary stenosis, ventricular septal defect, dextroposition of the aorta (aorta is on the right side instead of the left) and hypertrophy of the right ventricle. This condition results in a blue baby at birth due to inadequate oxygenation. Surgical

**Nanjing BYabscience technology Co.,Ltd**



correction is emergent.,domain:The CCHC-type zinc fingers 1, 5, 6 and 8 directly bind to GATA-type zinc fingers. The Tyr residue adjacent to the last Cys of the CCHC-type zinc finger is essential for the interact

## Background

zinc finger protein, FOG family member 2(ZFPM2) Homo sapiens The zinc finger protein encoded by this gene is a widely expressed member of the FOG family of transcription factors. The family members modulate the activity of GATA family proteins, which are important regulators of hematopoiesis and cardiogenesis in mammals. It has been demonstrated that the protein can both activate and down-regulate expression of GATA-target genes, suggesting different modulation in different promoter contexts. A related mRNA suggests an alternatively spliced product but this information is not yet fully supported by the sequence. [provided by RefSeq, Jul 2008],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using FOG-2 Monoclonal Antibody

**Nanjing BYabs science technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658