



IL-17RB Monoclonal Antibody

Catalog No	BYmab-10598
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	IL17RB
Protein Name	IL-17RB
Immunogen	Synthesized peptide derived from IL-17RB at AA range: 251-300
Specificity	IL-17RB Monoclonal Antibody detects endogenous levels of IL-17RB
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Interleukin-17 receptor B (IL-17 receptor B) (IL-17RB) (Cytokine receptor-like 4) (IL-17 receptor homolog 1) (IL-17Rh1) (IL17Rh1) (Interleukin-17B receptor) (IL-17B receptor)
Observed Band	55kD
Cell Pathway	[Isoform 1]: Cell membrane; Single-pass type I membrane protein.; [Isoform 2]: Secreted.
Tissue Specificity	Expressed in several endocrine tissues, mostly in fetal and adult liver, kidney, pancreas, testis, colon, brain and small intestine; not detected in peripheral blood leukocytes, lymphoid organs, and most cell lines.
Function	function:Receptor for the proinflammatory cytokines IL17B and IL17E. May play a role in controlling the growth and/or differentiation of hematopoietic cells.,tissue specificity:Expressed in several endocrine tissues, mostly in fetal and adult liver, kidney, pancreas, testis, colon, brain and small intestine; not detected in peripheral blood leukocytes, lymphoid organs, and most cell lines.,
Background	The protein encoded by this gene is a cytokine receptor. This receptor specifically binds to IL17B and IL17E, but does not bind to IL17 and IL17C. This

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务,欢迎咨询

	receptor has been shown to mediate the activation of NF-kapMAB and the production of IL8 induced by IL17E. The expression of the rat counterpart of this gene was found to be significantly up-regulated during intestinal inflammation, which suggested the immunoregulatory activity of this receptor. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using IL-17RB Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd