



# CYFIP2 Monoclonal Antibody

<b>Catalog No</b>	BYmab-10599
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	CYFIP2
<b>Protein Name</b>	CYFIP2
<b>Immunogen</b>	Synthesized peptide derived from CYFIP2 at AA range: 1171-1220
<b>Specificity</b>	CYFIP2 Monoclonal Antibody detects endogenous levels of CYFIP2
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Cytoplasmic FMR1-interacting protein 2 (p53-inducible protein 121)
<b>Observed Band</b>	150 45kD
<b>Cell Pathway</b>	Cytoplasm . Nucleus . Cytoplasm, perinuclear region . Cell junction, synapse, synaptosome . Highly expressed in the perinuclear regionand enriched in synaptosomes (By similarity). Treatment with leptomycin-B triggers translocation to the nucleus (PubMed:17245118). .
<b>Tissue Specificity</b>	Expressed in T-cells. Increased expression is observed in CD4(+) T-lymphocytes from patients with multiple sclerosis (at protein level).
<b>Function</b>	disease:Up-regulated significantly in CD4+ T lymphocytes from patients with multiple sclerosis (at protein level).,function:Involved in T-cell adhesion and p53-dependent induction of apoptosis. Does not bind RNA.,induction:By p53.,RNA editing:Partially edited. Editing appears to be brain-specific.,similarity:Belongs to the CYFIP family.,subcellular location:Highly expressed in the perinuclear region. Enriched in synaptosomes. Treatment with leptomycin-B triggers translocation to the nucleus.,subunit:Interacts with FMR1, FXR1 AND FXR2. Component of the WAVE1 complex composed of ABI2, CYFIP2, C3orf10/HSPC300, NCKAP1 and WASF1/WAVE1. CYFIP2 binds to

**Nanjing BYabscience technology Co.,Ltd**



activated RAC1 which causes the complex to dissociate, releasing activated WASF1. The complex can also be activated by NCK1.,

## Background

disease:Up-regulated significantly in CD4+ T lymphocytes from patients with multiple sclerosis (at protein level).,function:Involved in T-cell adhesion and p53-dependent induction of apoptosis. Does not bind RNA.,induction:By p53.,RNA editing:Partially edited. Editing appears to be brain-specific.,similarity:Belongs to the CYFIP family.,subcellular location:Highly expressed in the perinuclear region. Enriched in synaptosomes. Treatment with leptomycin-B triggers translocation to the nucleus.,subunit:Interacts with FMR1, FXR1 AND FXR2. Component of the WAVE1 complex composed of ABI2, CYFIP2, C3orf10/HSPC300, NCKAP1 and WASF1/WAVE1. CYFIP2 binds to activated RAC1 which causes the complex to dissociate, releasing activated WASF1. The complex can also be activated by NCK1.,


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using CYFIP2 Monoclonal Antibody