



# TyrRS Monoclonal Antibody

Catalog No	BYmab-10608
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	YARS
Protein Name	TyrRS
Immunogen	The antiserum was produced against synthesized peptide derived from the C-terminal region of human YARS. AA range:451-500
Specificity	TyrRS Monoclonal Antibody detects endogenous levels of TyrRS
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Tyrosine--tRNA ligase, cytoplasmic (EC 6.1.1.1) (Tyrosyl-tRNA synthetase) (TyrRS)
Observed Band	60kD
Cell Pathway	Cytoplasm .
Tissue Specificity	B-cell lymphoma,Bone,Lung,Platelet,
Function	catalytic activity:ATP + L-tyrosine + tRNA(Tyr) = AMP + diphosphate + L-tyrosyl-tRNA(Tyr).,disease:Defects in YARS are the cause of Charcot-Marie-Tooth disease dominant intermediate type C (CMTDIC) [MIM:608323]. CMTDIC is a form of Charcot-Marie-Tooth disease characterized by clinical and pathologic features intermediate between demyelinating and axonal peripheral neuropathies, and motor median nerve conduction velocities ranging from 25 to 45 m/sec.,function:Catalyzes the attachment of tyrosine to tRNA(Tyr) in a two-step reaction: tyrosine is first activated by ATP to form Tyr-AMP and then transferred to the acceptor end of tRNA(Tyr).,similarity:Belongs to the class-I aminoacyl-tRNA synthetase

**Nanjing BYabscience technology Co.,Ltd**



family.,similarity:Contains 1 tRNA-binding domain.,subunit:Homodimer.,

## Background

Aminoacyl-tRNA synthetases catalyze the aminoacylation of tRNA by their cognate amino acid. Because of their central role in linking amino acids with nucleotide triplets contained in tRNAs, aminoacyl-tRNA synthetases are thought to be among the first proteins that appeared in evolution. Tyrosyl-tRNA synthetase belongs to the class I tRNA synthetase family. Cytokine activities have also been observed for the human tyrosyl-tRNA synthetase, after it is split into two parts, an N-terminal fragment that harbors the catalytic site and a C-terminal fragment found only in the mammalian enzyme. The N-terminal fragment is an interleukin-8-like cytokine, whereas the released C-terminal fragment is an EMAP II-like cytokine. [provided by RefSeq, Jul 2008],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using TyrRS Monoclonal Antibody

**Nanjing BYabs science technology Co.,Ltd**

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658