



## FAM48A Monoclonal Antibody

BYmab-10628IgGHuman;Mouse;RatWBFAM48A C13orf19 FP757family with sequence similarity 48, member AThe antiserum was produced against synthesized peptide derived from the Internal region of human FAM48A. AA range:231-280The antibody detects endogenous FAM48A proteinLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Monoclonal, Mouse,IgGThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.WB 1:500-20001 mg/ml
Human;Mouse;Rat   WB   FAM48A C13orf19 FP757   family with sequence similarity 48, member A   The antiserum was produced against synthesized peptide derived from the Internal region of human FAM48A. AA range:231-280   The antibody detects endogenous FAM48A protein   Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   Monoclonal, Mouse,IgG   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   WB 1:500-2000
WBFAM48A C13orf19 FP757family with sequence similarity 48, member AThe antiserum was produced against synthesized peptide derived from the Internal region of human FAM48A. AA range:231-280The antibody detects endogenous FAM48A proteinLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Monoclonal, Mouse,IgGThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.WB 1:500-2000
FAM48A C13orf19 FP757family with sequence similarity 48, member AThe antiserum was produced against synthesized peptide derived from the Internal region of human FAM48A. AA range:231-280The antibody detects endogenous FAM48A proteinLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Monoclonal, Mouse,IgGThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.WB 1:500-2000
family with sequence similarity 48, member AThe antiserum was produced against synthesized peptide derived from the Internal region of human FAM48A. AA range:231-280The antibody detects endogenous FAM48A proteinLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Monoclonal, Mouse,IgGThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.WB 1:500-2000
The antiserum was produced against synthesized peptide derived from the Internal region of human FAM48A. AA range:231-280The antibody detects endogenous FAM48A proteinLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Monoclonal, Mouse,IgGThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.WB 1:500-2000
Internal region of human FAM48A. AA range:231-280 The antibody detects endogenous FAM48A protein Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Monoclonal, Mouse,IgG The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. WB 1:500-2000
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Monoclonal, Mouse,IgG The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. WB 1:500-2000
Monoclonal, Mouse,IgG The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. WB 1:500-2000
The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. WB 1:500-2000
affinity-chromatography using epitope-specific immunogen. WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
FAM48A C13orf19 FP757
85kD
Nucleus .
Highly expressed in testis, moderately in brain and pituitary gland. Expressed in several fetal tissues, including lung, brain, thymus and kidney. Expression is down-regulated in malignant prostate tissues.
function:Required for MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) activation during gastrulation. Required for down-regulation of E-cadherin during gastrulation by regulating E-cadherin protein level downstream from NCK-interacting kinase (NIK) and independently of the regulation of transcription by Fgf signaling and Snail.,similarity:Belongs to the FAM48 family.,subunit:Interacts with MAPK14.,tissue specificity:Highly expressed in testis, moderately in brain and pituitary gland. Expressed in several fetal tissues, including lung, brain, thymus and kidney. Expression is down-regulated in

## Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	function:Required for MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) activation during gastrulation. Required for down-regulation of E-cadherin during gastrulation by regulating E-cadherin protein level downstream from NCK-interacting kinase (NIK) and independently of the regulation of transcription by Fgf signaling and Snail.,similarity:Belongs to the FAM48 family.,subunit:Interacts with MAPK14.,tissue specificity:Highly expressed in testis, moderately in brain and pituitary gland. Expressed in several fetal tissues, including lung, brain, thymus and kidney. Expression is down-regulated in malignant prostate tissues.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

Western Blot analysis of various cells using FAM48A Monoclonal Antibody



Nanjing BYabscience technology Co.,Ltd