



PAR1 (Cleaved-Ser42) Monoclonal Antibody

Catalog No	BYmab-10822
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	F2R CF2R PAR1 TR
Protein Name	Proteinase-activated receptor 1 (PAR-1) (Coagulation factor II receptor) (Thrombin receptor)
Immunogen	Synthesized peptide derived from human PAR1 (Cleaved-Ser42) Monoclonal
Specificity	This antibody detects endogenous levels of PAR1 (Cleaved-Ser42).
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year Proteinase-activated receptor 1 (PAR-1) (Coagulation factor II receptor)
Purity Storage Stability Synonyms	≥90% -20°C/1 year Proteinase-activated receptor 1 (PAR-1) (Coagulation factor II receptor) (Thrombin receptor)
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Proteinase-activated receptor 1 (PAR-1) (Coagulation factor II receptor) (Thrombin receptor) 46kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Proteinase-activated receptor 1 (PAR-1) (Coagulation factor II receptor) (Thrombin receptor) 46kD Cell membrane; Multi-pass membrane protein.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



splicing results in multiple transcript variants. [provided by RefSeq, Aug 2015],

matters	needing
attentio	

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images Western Blot analysis of various cells using PAR1 (Cleaved-Ser42) Monoclonal Antibody 15-