



IDHP Monoclonal Antibody

Catalog No	BYmab-10826
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	IDH2
Protein Name	IDHP
Immunogen	Synthesized peptide derived from human IDHP
Specificity	This antibody detects endogenous levels of human IDHP
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Isocitrate dehydrogenase [NADP], mitochondrial (IDH;EC 1.1.1.42;ICD-M;IDP;NADP(+)-specific ICDH;Oxalosuccinate decarboxylase)
Observed Band	48kD
Cell Pathway	Mitochondrion .
Tissue Specificity	Colon,Heart,Kidney,Skin,
Function	catalytic activity:Isocitrate + NADP(+) = 2-oxoglutarate + CO(2) + NADPH.,catalytic activity:Oxalosuccinate + NADP(+) = 2-oxoglutarate + CO(2) + NADPH.,cofactor:Binds 1 magnesium or manganese ion per subunit.,function:Plays a role in intermediary metabolism and energy production. It may tightly associate or interact with the pyruvate dehydrogenase complex.,similarity:Belongs to the isocitrate and isopropylmalate dehydrogenases family.,subunit:Homodimer.,
Background	Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate

Nanjing BYabscience technology Co.,Ltd



dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydrogenase found in the mitochondria. It plays a role in intermediary metabolism and energy production. This protein may tightly associate or interact with the pyruvate dehydrogenase complex. Alternative splicing results in multiple transcript variants. [provided by Ref

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images