



TCEAL7 Polyclona Antibody

Isotype IgG Reactivity Human; Mouse Applications WB Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7; Transcription elongation factor S-II protein-like 7) Observed Band . Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function Turetus and ovary. Down-regulated in epithelial ovarian, tumors.<	Isotype IgG Reactivity Human; Mouse Applications WB Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uters and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breas brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation, similarity:Belongs to the TFS-II family. TFA subfamily., <t< th=""><th></th><th></th></t<>		
Reactivity Human; Mouse Applications WB Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Ucleus . Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Reactivity Human; Mouse Applications WB Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Uuesus . Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TF	Catalog No	BYmab-10892
Applications WB Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Lifely expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily.	Applications WB Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Tissue Specificity Flinction Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung carcer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily.,	lsotype	lgG
Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Nucleus . Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.	Gene Name TCEAL7 Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcriptio elongation factor S-II protein-like 7) Observed Band Uncleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breasi brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily.,	Reactivity	Human; Mouse
Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.	Protein Name TCEAL7 Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcriptio elongation factor S-II protein-like 7) Observed Band Uucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breasi brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Applications	WB
Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Liquid y expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family.	Immunogen Synthesized peptide derived from human TCEAL7 AA range: 34-84 Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcriptio elongation factor S-II protein-like 7) Observed Band Uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breasi brain and lung cancer cell lines and in brain and ovarian, cervical, prostate, breasi brain and lung cancer cell lines and in brain and ovariant tumors. Function function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family.	Gene Name	TCEAL7
Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the	Specificity This antibody detects endogenous levels of human TCEAL7 Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breasi brain and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breasi brain and ovary. Down-regulated in epithelial ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Protein Name	TCEAL7
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Ucleus . Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Eel Pathway Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breasi brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Immunogen	Synthesized peptide derived from human TCEAL7 AA range: 34-84
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Nucleus . Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breasi brain and lung cancer cell lines and in brain and ovarian tumors. Function function: May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily., Background function: May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily.,	Specificity	This antibody detects endogenous levels of human TCEAL7
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the	Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the	Purity ≥90% Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability-20°C/1 yearSynonymsTranscription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7)Observed BandCell PathwayCell PathwayNucleus .Tissue SpecificityHighly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the	Storage Stability -20°C/1 year Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Cell Pathway Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
SynonymsTranscription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7)Observed BandNucleus .Cell PathwayNucleus .Tissue SpecificityHighly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the	Synonyms Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7) Observed Band Nucleus . Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation., similarity:Belongs to the TFS-II family. TFA subfamily., matters needing Avoid repeated freezing and thawing!	Purity	≥90%
elongation factor S-II protein-like 7)Observed BandCell PathwayNucleus .Tissue SpecificityHighly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the	elongation factor S-II protein-like 7) Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors. Function function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell PathwayNucleus .Tissue SpecificityHighly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the	Cell PathwayNucleus .Tissue SpecificityHighly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,matters needingAvoid repeated freezing and thawing!	Synonyms	Transcription elongation factor A protein-like 7 (TCEA-like protein 7;Transcription elongation factor S-II protein-like 7)
Tissue SpecificityHighly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the	Tissue SpecificityHighly expressed in normal and fetal brain tissues, and weakly expressed in uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,matters needingAvoid repeated freezing and thawing!	Observed Band	
uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast, brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the	uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast brain and lung cancer cell lines and in brain and ovarian tumors.Functionfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,matters needingAvoid repeated freezing and thawing!	Cell Pathway	Nucleus .
TFS-II family. TFA subfamily.,Backgroundfunction:May be involved in transcriptional regulation., similarity:Belongs to the	TFS-II family. TFA subfamily., Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	uterus and ovary. Down-regulated in epithelial ovarian, cervical, prostate, breast,
Background function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,	TFS-II family. TFA subfamily., matters needing Avoid repeated freezing and thawing!	Function	function:May be involved in transcriptional regulation.,similarity:Belongs to the TFS-II family. TFA subfamily.,
		Background	function:May be involved in transcriptional regulation.,similarity:Belongs to the
matters needing Avoid repeated freezing and thawing! attention		matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TCEAL7 Polyclona Antibody

Nanjing BYabscience technology Co.,Ltd