





OSBL8 mouse mAb

| Catalog No | BYmab-11044 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | OSBPL8 KIAA1451 ORP8 OSBP10 |
| Protein Name | OSBL8 |
| Immunogen | Synthesized peptide derived from human OSBL8 AA range: 290-340 |
| Specificity | This antibody detects endogenous levels of OSBL8 at Human |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| Cell Pathway | [Isoform 1]: Endoplasmic reticulum membrane; Single-pass membrane protein. Nucleus membrane. The presence of the N-terminus extension contains an overall negative charge that may explain the weak localization to the cortical endoplasmic reticulum (Probable); [Isoform 3]: Endoplasmic reticulum membrane; Single-pass membrane protein. Localizes to the cortical endoplasmic reticulum at the endoplasmic reticulum-plasma membrane contact sites |
| Tissue Specificity | Widely expressed (PubMed:11735225). Expressed at higher level in macrophages (PubMed:17991739). |
| Function | similarity:Belongs to the OSBP family.,similarity:Contains 1 PH domain.,tissue specificity:Widely expressed., |
| Background | This gene encodes a member of a family of proteins containing an N-terminal pleckstrin homology domain and a highly conserved C-terminal oxysterol-binding protein-like sterol-binding domain. It binds mutliple lipid-containing molecules, including phosphatidylserine, phosphatidylinositol 4-phosphate (PI4P) and oxysterol, and promotes their exchange between the endoplasmic reticulum and |
| | Nanjing Pyahosianos tachnalogy Co. Ltd. |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| | the plasma membrane. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |



