



FBX41 mouse mAb

Catalog No BYmab-11129 Isotype IgG Reactivity Human; Mouse Applications WB Gene Name FBXO41 FBX41 KIAA1940 Protein Name FBX41 Immunogen Synthesized peptide derived from human FBX41 AA range: 622-672 Specificity This antibody detects endogenous levels of FBX41 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function: Substrate-recognition component of the SCF (SKP1-CUL1-F-box domain, subunit:Directly interacts with SKP1A and CUL1.1 Background This gene encodes a member of the F-box protein inquise complex that plays a role in phosphorylation-dependent ubiquitination. F-box protein is a role in phosphorylation-dependent ubiquitination. F-box proteins are obtained in addition to the		
Reactivity Human; Mouse Applications WB Gene Name FBX041 FBX41 KIAA1940 Protein Name FBX41 Immunogen Synthesized peptide derived from human FBX41 AA range: 622-672 Specificity This antibody detects endogenous levels of FBX41 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function: Substrate-recognition component of the SCF (SKP1-CUL1.F-box protein)-type E3 ubiquitin ligase complex, similarity. Contains 1 F-box domain, subunit: Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif. FBXV proteins contain in liter different	Catalog No	BYmab-11129
Applications Gene Name FBXO41 FBX41 KIAA1940 Protein Name FBX41 Immunogen Synthesized peptide derived from human FBX41 AA range: 622-672 Specificity This antibody detects endogenous levels of FBX41 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function: Substrate-recognition component of the SCF (SKP1-CUL1.F-box protein)-type E3 ubiquitin ligase complex, similarity: Contains 1 F-box domain, subunit: Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box, F-box proteins constitute one classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXV proteins contain either different	Isotype	IgG
Gene Name FBXO41 FBX41 KIAA1940 Protein Name FBX41 Immunogen Synthesized peptide derived from human FBX41 AA range: 622-672 Specificity This antibody detects endogenous levels of FBX41 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex, similarity. Contains 1 F-box domain., subunit:Directly inferacts with SKP1A and CUL1. Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute on of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 doma	Reactivity	Human; Mouse
Protein Name FBX41 Immunogen Synthesized peptide derived from human FBX41 AA range: 622-672 Specificity This antibody detects endogenous levels of FBX41 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex, similarity:Contains 1 F-box domain, subunit:Directly inferacts with SKP1A and CUL1-, Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid molif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXD proteins are divided into t	Applications	WB
Immunogen Synthesized peptide derived from human FBX41 AA range: 622-672 Specificity This antibody detects endogenous levels of FBX41 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex, similarity:Contains 1 F-box domain, subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contain in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain in deliction in addition to the F-box motif: repeats, and FBXD proteins can each contain either different	Gene Name	FBXO41 FBX41 KIAA1940
Specificity This antibody detects endogenous levels of FBX41 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex, similarity:Contains 1 F-box domain, subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Protein Name	FBX41
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function: Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex., similarity: Contains 1 F-box domain., subunit: Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain wD-40 domains, FBXL proteins contain leuteni-rich repeats, and FBXD proteins contain either different	Immunogen	Synthesized peptide derived from human FBX41 AA range: 622-672
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex, similarity:Contains 1 F-box domain., subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain elution trick represents and EBXV proteins contain in addition to the F-box motif: FBXW proteins contain in addition to the F-box motif: FBXW proteins contain in addition to the fifteent ifferent	Specificity	This antibody detects endogenous levels of FBX41 at Human/Mouse
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complexsimilarity:Contains 1 F-box domainsubunit:Directly interacts with SKP1A and CUL1 Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain wD-40 domains, FBXL proteins contain letther different	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity.Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Source	Monoclonal, Mouse,IgG
Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex_similarity:Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity:Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Dilution	
Synonyms Observed Band Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity:Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Tissue Specificity Function function: Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity: Contains 1 F-box domain.,subunit: Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Purity	≥90%
Cell Pathway Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity:Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Storage Stability	-20°C/1 year
Tissue Specificity Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity:Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Synonyms	
Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity:Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., Background This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different		
Function function:Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex.,similarity:Contains 1 F-box domain.,subunit:Directly interacts with SKP1A and CUL1., This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Observed Band	
protein)-type E3 ubiquitin ligase complex., similarity: Contains 1 F-box domain., subunit: Directly interacts with SKP1A and CUL1., This gene encodes a member of the F-box protein family, which is characterized by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different		
by an approximately 40 amino acid motif, the F-box. F-box proteins constitute one of the four subunits of the SCF ubiquitin protein ligase complex that plays a role in phosphorylation-dependent ubiquitination. F-box proteins are divided into three classes depending on the interaction substrate domain each contains in addition to the F-box motif: FBXW proteins contain WD-40 domains, FBXL proteins contain leucine-rich repeats, and FBXO proteins contain either different	Cell Pathway	
	Cell Pathway Tissue Specificity	protein)-type E3 ubiquitin ligase complex.,similarity:Contains 1 F-box


Nanjing BYabscience technology Co.,Ltd





	encoded by this gene belongs to the FBXO class. [provided by RefSeq, Feb 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658