



## NIT2 mouse mAb

Catalog No	BYmab-11174
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	NIT2 CUA002
Protein Name	NIT2
Immunogen	Synthesized peptide derived from human NIT2 AA range: 200-250
Specificity	This antibody detects endogenous levels of NIT2 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	<del>-</del>
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Cytoplasm .  Detected in fetal brain (at protein level). Ubiquitous. Detected in heart, brain,
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  Cytoplasm .  Detected in fetal brain (at protein level). Ubiquitous. Detected in heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, prostate, spleen, thymus, prostate, testis, ovary, small intestine and colon.  function:Overexpression decreases the colony-forming capacity of cultured cells by arresting cells in the G2 phase of the cell cycle.,mass spectrometry: PubMed:17488281,similarity:Belongs to the UPF0012 family.,similarity:Contains 1 CN hydrolase domain.,tissue specificity:Detected in fetal brain (at protein level).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Cytoplasm .  Detected in fetal brain (at protein level). Ubiquitous. Detected in heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, prostate, spleen, thymus, prostate, testis, ovary, small intestine and colon.  function:Overexpression decreases the colony-forming capacity of cultured cells by arresting cells in the G2 phase of the cell cycle.,mass spectrometry: PubMed:17488281,similarity:Belongs to the UPF0012 family.,similarity:Contains 1 CN hydrolase domain.,tissue specificity:Detected in fetal brain (at protein level). Ubiquitous. Detected in heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, prostate, spleen, thymus, prostate, testis, ovary, small intestine and

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

