





F8I2 mouse mAb

Catalog No BYmab-11229 Isotype IgG Reactivity Human; Mouse Applications WB Gene Name F8A1 F8A; F8A2 F8A; F8A3 F8A Protein Name F8I2 Immunogen Synthesized peptide derived from human F8I2 AA range: 33-83 Specificity This antibody detects endogenous levels of F8I2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In euron found in Intranuclear structures, the intranuclear rodles (RVRs), also known as rodlets of Roncoroni, in association with theighting (Mgs) milarity. Collocalized with HTT in endosomes (PubMed:164783 in encorn found in Intranuclear structures, the intranuclear rodles (RVRs), also known as rodlets of Roncoroni, in associ		
Reactivity Applications WB Gene Name F8A1 F8A; F8A2 F8A; F8A3 F8A Protein Name F8I2 Immunogen Synthesized peptide derived from human F8I2 AA range: 33-83 Specificity This antibody detects endogenous levels of F8I2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm , Nucleus . Early endosome , Nucleus, nuclear body , Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed: 16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed: 6476778) In neuron found in intranuclear structures, the intranuclear roddets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Catalog No	BYmab-11229
Applications Gene Name F8A1 F8A; F8A2 F8A; F8A3 F8A Protein Name F8I2 Immunogen Synthesized peptide derived from human F8I2 AA range: 33-83 Specificity This antibody detects endogenous levels of F8I2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed: 16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed: 6476778) In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function In function: Not known. Possible housekeeping role, tissue specificity: Produced abundantly in a wide variety of cell types. Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (inti22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Isotype	IgG
Gene Name F8A1 F8A; F8A2 F8A; F8A3 F8A Protein Name F8I2 Immunogen Synthesized peptide derived from human F8I2 AA range: 33-83 Specificity This antibody detects endogenous levels of F8I2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT., F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778), in neuron found in intranuclear structures, the intranuclear roddets (INFs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role, tissue specificity:Produced abundantly in a wide variety of cell types. Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Reactivity	Human; Mouse
Protein Name F8I2 Immunogen Synthesized peptide derived from human F8I2 AA range: 33-83 Specificity This antibody detects endogenous levels of F8I2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cytoplasm Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role, tissue specificity: Produced abundantly in a wide variety of cell types. Background This gene is contained entirely within intron 22 of the factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to t	Applications	WB
Immunogen Synthesized peptide derived from human F8I2 AA range: 33-83 Specificity This antibody detects endogenous levels of F8I2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In neuron found in absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity) Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., Function This gene is contained entirely within intron 22 of the factor VIII, A portion of intron 22 (int22h), containing F8A, is repeated twice e	Gene Name	F8A1 F8A; F8A2 F8A; F8A3 F8A
Specificity This antibody detects endogenous levels of F8I2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm Nucleus . Early endosome Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRS), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types. Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (intro), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Protein Name	F8I2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the X telomere. Although its function is unknown, the observation that this gene is	Immunogen	Synthesized peptide derived from human F8I2 AA range: 33-83
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT. F8A1/F8A2/F8A3 is concentral in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRS), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xg telomere. Although its function is unknown, the observation that this gene is	Specificity	This antibody detects endogenous levels of F8I2 at Human/Mouse
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred (in cytoplasm (lNs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types. Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear redlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types. Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Ky telomere. Although its function is unknown, the observation that this gene is	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity) Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity) Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Purity	≥90%
Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity) Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Nucleus . Early endosome . Nucleus, nuclear body . Diffuse presence in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Synonyms	
in the cytoplasm and accumulation in the nucleus (PubMed:16476778). In absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of Roncoroni, in association with ubiquitin (By similarity). Tissue Specificity Produced abundantly in a wide variety of cell types. Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Observed Band	
Function function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types., This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Cell Pathway	absence of HTT, F8A1/F8A2/F8A3 is concentred in cytoplasm (By similarity). Colocalized with HTT in endosomes (PubMed:16476778). In neuron found in intranuclear structures, the intranuclear rodlets (INRs), also known as rodlets of
Background This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Tissue Specificity	Produced abundantly in a wide variety of cell types.
than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is	Function	function:Not known. Possible housekeeping role.,tissue specificity:Produced abundantly in a wide variety of cell types.,
	Background	than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	is transcribed abundantly in a wide variety of cell types. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



