



P4HA3 mouse mAb

Catalog No	BYmab-11252
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	P4HA3 UNQ711/PRO1374
Protein Name	P4HA3
Immunogen	Synthesized peptide derived from human P4HA3 AA range: 211-261
Specificity	This antibody detects endogenous levels of P4HA3 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Endoplasmic reticulum lumen .
Tissue Specificity	Highly expressed in placenta, liver and fetal skin. Weakly expressed in fetal epiphyseal cartilage, fetal liver, fibroblast, lung and skeletal muscle. Expressed also in fibrous cap of carotid atherosclerotic lesions.
Function	catalytic activity:Procollagen L-proline + 2-oxoglutarate + O(2) = procollagen trans-4-hydroxy-L-proline + succinate + CO(2).,cofactor:Ascorbate.,cofactor: Binds 1 Fe(2+) ion per subunit.,function:Catalyzes the post-translational formation of 4-hydroxyproline in -Xaa-Pro-Gly- sequences in collagens and other proteins.,PTM:N-glycosylation plays no role in the catalytic activity.,similarity:Belongs to the P4HA family.,similarity:Contains 1 PKHD (prolyl/lysyl hydroxylase) domain.,similarity:Contains 1 TPR repeat.,subunit:Heterotetramer of two alpha-3 chains and two beta chains (the beta chain is the multi-functional PDI).,tissue specificity:Highly expressed in placenta, liver and fetal skin. Weakly expressed in fetal epiphyseal cartilage, fetal liver, fibroblast, lung and skeletal muscle.


Nanjing BYabscience technology Co.,Ltd



Expressed also in fibrous cap of carotid atherosclerotic lesions.,

Background	This gene encodes a component of prolyl 4-hydroxylase, a key enzyme in collagen synthesis composed of two identical alpha subunits and two beta subunits. The encoded protein is one of several different types of alpha subunits and provides the major part of the catalytic site of the active enzyme. In collagen and related proteins, prolyl 4-hydroxylase catalyzes the formation of 4-hydroxyproline that is essential to the proper three-dimensional folding of newly synthesized procollagen chains. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2014],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using P4HA3 mouse mAb

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658