



MTDC mouse mAb

Catalog No	BYmab-11332
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	MTHFD2 NMDMC
Protein Name	MTDC
Immunogen	Synthesized peptide derived from human MTDC AA range: 121-171
Specificity	This antibody detects endogenous levels of MTDC at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Mitochondrion.
Tissue Specificity	
Function	catalytic activity:5,10-methenyltetrahydrofolate + H(2)O = 10-formyltetrahydrofolate.,catalytic activity:5,10-methylenetetrahydrofolate + NAD(+) = 5,10-methenyltetrahydrofolate + NADH.,cofactor:Magnesium.,developmental stage:Expressed only in developing normal tissues.,miscellaneous:This NAD-dependent bifunctional enzyme has very different kinetic properties than the larger NADP-dependent trifunctional enzyme and is unique in that it requires formation of an enzyme-magnesium complex to allow binding of NAD.,similarity:Belongs to the tetrahydrofolate dehydrogenase/cyclohydrolase family.,subunit:Homodimer.,
Background	This gene encodes a nuclear-encoded mitochondrial bifunctional enzyme with methylenetetrahydrofolate dehydrogenase and methenyltetrahydrofolate

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	cyclohydrolase activities. The enzyme functions as a homodimer and is unique in its absolute requirement for magnesium and inorganic phosphate. Formation of the enzyme-magnesium complex allows binding of NAD. Alternative splicing results in two different transcripts, one protein-coding and the other not protein-coding. This gene has a pseudogene on chromosome 7. [provided by RefSeq, Mar 2009],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For

Products Images



Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658