



APEX2 mouse mAb

Catalog No	BYmab-11334
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	APEX2 APE2 APEXL2 XTH2
Protein Name	APEX2
Immunogen	Synthesized peptide derived from human APEX2 AA range: 196-246
Specificity	This antibody detects endogenous levels of APEX2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus. Cytoplasm. Mitochondrion . Together with PCNA, is redistributed in discrete nuclear foci in presence of oxidative DNA damaging agents.
Tissue Specificity	Highly expressed in brain and kidney. Weakly expressed in the fetal brain.
Function	catalytic activity:The C-O-P bond 3' to the apurinic or apyrimidinic site in DNA is broken by a beta-elimination reaction, leaving a 3'-terminal unsaturated sugar and a product with a terminal 5'-phosphate.,function:May participate in both nuclear and mitochondrial post-replicative base excision repair (BER). In the nucleus functions in the PCNA-dependent BER pathway.,similarity:Belongs to the DNA repair enzymes AP/exoA family.,subcellular location:Colocalized partly with PCNA in nuclear foci.,subunit:Interacts with PCNA. This interaction is increased by misincorporation of uracil in nuclear DNA.,tissue specificity:Highly expressed in cells, adult brain and kidney. Weakly expressed in the fetal brain.,
Background	Apurinic/apyrimidinic (AP) sites occur frequently in DNA molecules by spontaneous hydrolysis, by DNA damaging agents or by DNA glycosylases that

Nanjing BYabscience technology Co.,Ltd



remove specific abnormal bases. AP sites are pre-mutagenic lesions that can prevent normal DNA replication so the cell contains systems to identify and repair such sites. Class II AP endonucleases cleave the phosphodiester backbone 5' to the AP site. This gene encodes a protein shown to have a weak class II AP endonuclease activity. Most of the encoded protein is located in the nucleus but some is also present in mitochondria. This protein may play an important role in both nuclear and mitochondrial base excision repair. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2012],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using APEX2 mouse mAb