



## LOXL4 mouse mAb

Catalog No	BYmab-11391
Isotype	lgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	LOXL4 LOXC
Protein Name	LOXL4
Immunogen	Synthesized peptide derived from human LOXL4 AA range: 39-89
Specificity	This antibody detects endogenous levels of LOXL4 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Secreted, extracellular space .
Tissue Specificity	Expressed in many tissues, the highest levels among the tissues studied being in the skeletal muscle, testis and pancreas. Expressed in cartilage.
Function	cofactor:Contains 1 lysine tyrosylquinone.,cofactor:Copper.,function:May modulate the formation of a collagenous extracellular matrix.,PTM:The lysine tyrosylquinone cross-link (LTQ) is generated by condensation of the epsilon-amino group of a lysine with a topaquinone produced by oxidation of tyrosine.,similarity:Belongs to the lysyl oxidase family.,similarity:Contains 4 SRCR domains.,tissue specificity:Expressed in many tissues, the highest levels among the tissues studied being in the skeletal muscle, testis and pancreas. Expressed in cartilage.,
Background	This gene encodes a member of the lysyl oxidase gene family. The prototypic member of the family is essential to the biogenesis of connective tissue, encoding an extracellular copper-dependent amine oxidase that catalyses the first step in
	Nanjing BYabscience technology Co.,Ltd



45 --

35 --

25 ---

15 --

10 ---

国内优质抗体供应商

精准的 WB 检测服务 (2)。

24H 在线服务,欢迎咨询



	24日任线服务,从应答问 国际关键	
	the formation of crosslinks in collagens and elastin. A highly conserved amino acid sequence at the C-terminus end appears to be sufficient for amine oxidase activity, suggesting that each family member may retain this function. The N-terminus is poorly conserved and may impart additional roles in developmental regulation, senescence, tumor suppression, cell growth control, and chemotaxis to each member of the family. [provided by RefSeq, Jul 2008],	
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	
Products Images		
1 kDa 180 140 100 75 60	Western Blot analysis of various cells using LOXL4 mouse mAb	

Nanjing BYabscience technology Co.,Ltd