



## PVRIG mouse mAb

Catalog No	BYmab-11627
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	PVRIG C7orf15
Protein Name	PVRIG
Immunogen	Synthesized peptide derived from human PVRIG AA range: 229-279
Specificity	This antibody detects endogenous levels of PVRIG at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	
otorago otability	-20°C/1 year
Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year  Cell membrane ; Multi-pass membrane protein .
Synonyms Observed Band	Cell membrane; Multi-pass membrane protein.  Expressed in some types of immune cells. Expressed at low levels on the surface of freshly isolated T-cells and natural killer (NK) cells, predominantly on CD8+T-cells (mainly memory/effector, but not naive cells) and on both CD16+ and CD16- NK cells. T-cell expression levels are variable among individuals. Not
Synonyms Observed Band Cell Pathway	Cell membrane; Multi-pass membrane protein.  Expressed in some types of immune cells. Expressed at low levels on the surface of freshly isolated T-cells and natural killer (NK) cells, predominantly on CD8+ T-cells (mainly memory/effector, but not naive cells) and on both CD16+ and CD16- NK cells. T-cell expression levels are variable among individuals. Not detected in B-cells, naive or helper T-cells, monocytes, nor neutrophils (at protein level). Not detected in dendritic cells.  caution:Despite its name, it does not share sequence similarity with the poliovirus
Synonyms  Observed Band  Cell Pathway  Tissue Specificity	Cell membrane; Multi-pass membrane protein.  Expressed in some types of immune cells. Expressed at low levels on the surface of freshly isolated T-cells and natural killer (NK) cells, predominantly on CD8+ T-cells (mainly memory/effector, but not naive cells) and on both CD16+ and CD16- NK cells. T-cell expression levels are variable among individuals. Not detected in B-cells, naive or helper T-cells, monocytes, nor neutrophils (at protein level). Not detected in dendritic cells.
Synonyms  Observed Band  Cell Pathway  Tissue Specificity  Function	Cell membrane; Multi-pass membrane protein.  Expressed in some types of immune cells. Expressed at low levels on the surface of freshly isolated T-cells and natural killer (NK) cells, predominantly on CD8+ T-cells (mainly memory/effector, but not naive cells) and on both CD16+ and CD16- NK cells. T-cell expression levels are variable among individuals. Not detected in B-cells, naive or helper T-cells, monocytes, nor neutrophils (at protein level). Not detected in dendritic cells.  caution:Despite its name, it does not share sequence similarity with the poliovirus

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using PVRIG mouse mAb