



M21D1 mouse mAb

Catalog No	BYmab-11697
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	MB21D1 C6orf150
Protein Name	M21D1
Immunogen	Synthesized peptide derived from human M21D1
Specificity	This antibody detects endogenous levels of M21D1 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	21780DA
Cell Pathway	Nucleus . Chromosome . Cell membrane ; Peripheral membrane protein . Cytoplasm, cytosol . Mainly localizes in the nucleus, and at low level in the cytosol (PubMed:31808743, PubMed:31544964). On chromosomes, enriched on centromeric satellite and LINE DNA repeat elements (PubMed:30811988). Exported from the nucleus to the cytosol in a XPO1/CRM1 via the nuclear export signal in response to DNA stimulation (PubMed:33406424). Outside the nucleus, localizes at the cell membrane as a peripheral membrane protein in resting conditions: association to the cell membrane is mediated via binding to phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P2) (PubMed:30827685). Localization at the cell membrane is required to limit the recognition of self-DNA (PubMed:30827685). Following detection of double-stranded DNA (dsDNA), released from the cell membrane into the cytosol in order to signal (PubMed:30827685). Upon transfection with dsDNA forms punctate structures that co-localize with DNA and Beclin-1 (BECN1) (PubMed:26048138). Phosphorylation at Tyr-215 promotes cytosolic retention (PubMed:30356214). In response to translation stress, translocates to the cytosol and associates with collided ribosomes (PubMed:34111399). .; (Microbial infection) Upon infection

Nanjing BYabscience technology Co.,Ltd



with virulent M.tuberculosis forms aggregates with dsDNA, non-virulent bacteria (without the ESX-1 locus) do not form these aggregates (PubMed:26048138). .

Tissue Specificity Expressed in the monocytic cell line THP1.

Function

Background

matters needing attention Avoid repeated freezing and thawing!

Usage suggestions This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998

监督电话: 15950492658