



## KBP mouse mAb

expressed at intermediate level in all other adult tissues examined, as well as in		
Reactivity   Human; Mouse;Rat     Applications   WB     Gene Name   KIAA1279 KBP     Protein Name   KBP     Immunogen   Synthesized peptide derived from human KBP AA range: 542-592     Spacificity   This antibody detects endogenous levels of KBP at Human/Mouse/Rat     Formulation   Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.     Source   Monoclonal, Mouse,IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   -     Cell Pathway   Cytoplasm, cytoskeleton .     Tissue Specificity   Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in fetal liver and brain. Not expressed in all other adult lisuses examined, as well as in tetal liver and brain. Not expressed in all other adult lisuses specific brain regions examined, as well as in tetal liver and brain. Not expressed in all other adult lisuses expressed at intermediate level in all other adult lisuses examined, as well as in tetal liver and brain. Not expressed in heart, braind, organ/ testion, organ/ zeton of axonal microtubules, and axon	Catalog No	BYmab-11739
ApplicationsWBGene NameKIAA1279 KBPProtein NameKBPImmunogenSynthesized peptide derived from human KBP AA range: 542-592SpecificityThis antibody detects endogenous levels of KBP at Human/Mouse/RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCytoplasm, cytoskeleton .Tissue SpecificityHighy expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult itsues examined, as well as in feature disputibility expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult itsues examined, as well as in feature adult fransports by mental retradation and facial dysmprism, as well as phenotypes related by herd at lever and all specific brain regions examined facial dysorphism, as well as phenotypes related by herd at lever and all specific prain regions examined and and all specific train regions examined, as well as in feature disputes of the KFF1-brinding protein fraindy expressed at intermediate level in all other adult incortypes related by herd at lever and all specific brain regions examined, as well as in feature synonyms	Isotype	lgG
Gene Name   KIAA1279 KBP     Protein Name   KBP     Immunogen   Synthesized peptide derived from human KBP AA range: 542-592     Specificity   This antibody detects endogenous levels of KBP at Human/Mouse/Rat     Formulation   Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.     Source   Monoclonal, Mouse,IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cytoplasm, cytoskeleton .     Tissue Specificity   Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in hefal liver and brain. Not expressed in blood leukcoytes.     Function   disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) IMMM:G094601, GOSHS is characterized by microcephaly, mediate levyin sall or dra and cand cephage related to the Hirscharpung disease syndrome. Function: Regulates micro and and groppine and and cephage related to the Hirscharpung disease syndrome. Function: Regulates micro and controm well as an hefal liver and brain. Not expressed in hord relating and central nervous system development. Regulates mined. Moderate expressed at intermediate level in all other adul	Reactivity	Human; Mouse;Rat
Protein NameKBPImmunogenSynthesized peptide derived from human KBP AA range: 542-592SpecificityThis antibody detects endogenous levels of KBP at Human/Mouse/RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearCell PathwayCytoplasm, cytoskeleton .Tissue SpecificityHighly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.Functiondisease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysorphism, as well as phenotypes related to hircsprung disease syndrome, function:Required for organization of axonal microtubles, and axonal outgrowth and maintenance during peripheral and central nervous system development. Function: Required to rorganization of axonal ming/way, subunit.Interacts with KIF 18, tissue specific/brain regions examined. Moderate expressed at intermediate level in all other adult tissue sexamined. Moderate expressed or the KIF 1-binding protein family.subunit.Interacts with KIF 18, tissue specific brain regions examined. Moderate expressed at intermediate level in all other adult tora	Applications	WB
ImmunogenSynthesized peptide derived from human KBP AA range: 542-592SpecificityThis antibody detects endogenous levels of KBP at Human/Mouse/RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsObserved BandCytoplasm, cytoskeleton .Tissue SpecificityHighly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.Functiondisease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460], GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phototypes related to hirschsprung disease syndrome. Junction. Required for organization of axonal microtubules, and axonal outgrowth and maintenance during periperal and central nervous system development. Regulates microchordral transport by modulating KIFT Bm to activity. Similarity Belongs to the KIFT-binitg protein and increaded for organization of axonal microtubules, spinal cord and all specific brain mergions examined. Axonal outgrowth and maintenance during periperal and central nervous system development. Regulates microchordral transport by modulating KIFT Bm theory activity. Similarity. Belongs to the KIFT-binitg protein disease syndrome. Junction Regulates for an negloon syndrome (axonal outgrowth and maint	Gene Name	KIAA1279 KBP
Specificity This antibody detects endogenous levels of KBP at Human/Mouse/Rat   Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -   Cell Pathway Cytoplasm, cytoskeleton .   Tissue Specificity Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) (MIM:609460). GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as well as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) (MIM:609460). GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease syndrome, function: Required for organization of axonal microbubles, and axonal ourigowith and maintenance during peripheral and	Protein Name	KBP
FormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsColloserved BandCell PathwayCytoplasm, cytoskeleton .Tissue SpecificityHighly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult ttssue system development. Not expressed in blood leukocytes.Functiondisease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon synomem (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirscherprug disease synthecytive, is spinal cord and all specific or and microtubules, and axonal outgrowth and maintenance during peripheral and microtubules, and axonal o	Immunogen	Synthesized peptide derived from human KBP AA range: 542-592
SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCytoplasm, cytoskeleton .Cell PathwayCytoplasm, cytoskeleton .Tissue SpecificityHighly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.Functiondisease :Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) (MIM:609460). GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome, function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity. similarity:Belongs to the KIF1-binding protein family, subunit.interacts with KIF1B. tissues examined, as well as in heart, brain and in a sevenila specific brain microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity. similarity:Belongs to the KIF1-binding protein family, subunit.interacts with KIF1B. tissue specific brain regions examined, as well as in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as perior for brain heart, brain, ovary, testis, spinal cord and all specif	Specificity	This antibody detects endogenous levels of KBP at Human/Mouse/Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cytoplasm, cytoskeleton .   Tissue Specificity Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome. (Linction:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondiral transport by modulating KIF1B motor activity. similarity:Belongs to the KIF1-binding protein family, subunit:Interacts with KIF1B., tissue specificity is examined, as well as in teart in all other adult tissues examined, as well as in termediate level in all other adult centra nervous system development. Regulates mitochondiral transport by modulating KIF1B motor activity. Similarity:Belongs to the KIF1-binding protein family, subunit:Interacts with KIF1B., tissue sexamined, as well as in tearl, brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in termediate level in all other adult tissues examined. Socherace the expresse	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cytoplasm, cytoskeleton .   Cell Pathway Cytoplasm, cytoskeleton .   Tissue Specificity Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in fetal liver and brain. Not expressed in blood lieukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, microtubules, and axonal outgrowth and maintenance during peripheral and microtubules, and axonal outgrowth and maintenance during peripheral and microtubules, and axonal outgrowth and maintenance during peripheral and microtubules, spinal cord and all specific brain regions examined, as well as in fetal liver and brain. Not expressed in blood lieukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, microtubules, and axonal outgrowth and maintenance during peripheral and microtubules, and axonal outgrowth and maintenance during peripheral and microtubules, and axonal outgrowth and maintenance during peripheral and microtubules, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined. Moderate expressed at intermediate level in all ot	Source	Monoclonal, Mouse,IgG
Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearObserved Band-20°C/1 yearCell PathwayCytoplasm, cytoskeleton .Tissue SpecificityHighly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.Functiondisease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome. Regulates mitochondrial transport by modulating KIF1B motor activity. similarity:Belongs to the KIF1-binding protein familysubunit:Interacts with KIF1B.tissue specificity:Highly expressed at intermediate level in all other adult torain ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined. And expressed at intermediate level in all other adult tissues examined at an and anticrotubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activitysimilarity:Belongs to the KIF1-binding protein familysubunit:Interacts with KIF1B.tissue specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in	Purification	
Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band -20°C/1 year   Cell Pathway Cytoplasm, cytoskeleton .   Tissue Specificity Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome. Requilates mitochondrial transport by modulating KIF1B motor activity, similarity:Belongs to the KIF1-binding protein family, subunit:Interacts with KIF1B, tissue specific ty:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in reative expressed at intermediate level in all other adult tissues examined. Moderate expressed at intermediate level in all other adult tissues examined. Moderate expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain regions examined, as well as infetal liver and brain transport by modulating KIF1B motor activity. Similarity:Belongs to the KIF1-binding protein family. Subunit:Interacts with KIF1B, tissue specific brain regions examined, as well as in theart,	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band -20°C/1 year   Cell Pathway Cytoplasm, cytoskeleton .   Tissue Specificity Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome. function: Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity, similarity:Belongs to the KIF1-binding protein family, subunit:Interacts with KIF1B., tissue specificity:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in termediate level in all other adult tissues examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in termediate level in all other adult tissues examined. Moderate expressed at intermediate level in all other adult tissues examined. Moderate expressed at intermediate level in all other adult tissues examined. Moderate expressed at intermediate level in all other adult tissues examined. Moderate expressed at intermediate level in all other adult tissues examined. Moderate expressed at intermediate level in all other adult tissues	Concentration	1 mg/ml
Synonyms   Observed Band   Cell Pathway Cytoplasm, cytoskeleton .   Tissue Specificity Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome., function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity., similarity:Belongs to the KIF1-binding protein family., subunit:Interacts with KIF1B., tissue specificity:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in the specific brain regions examined.	Purity	≥90%
Observed Band Cytoplasm, cytoskeleton .   Tissue Specificity Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.   Function disease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome., function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity., similarity:Belongs to the KIF1-binding protein family., subunit:Interacts with KIF1B., tissue specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in	Storage Stability	-20°C/1 year
Cell PathwayCytoplasm, cytoskeleton .Tissue SpecificityHighly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.Functiondisease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome.,function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity.,similarity:Belongs to the KIF1-binding protein family.,subunit:Interacts with KIF1B.,tissue specificity:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in	Synonyms	
Tissue SpecificityHighly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood leukocytes.Functiondisease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome.,function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity.,similarity:Belongs to the KIF1-binding protein family.,subunit:Interacts with KIF1B.,tissue specificity:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in	Observed Band	
Functiondisease:Defects in KIAA1279 are the cause of Goldberg-Shprintzen megacolon syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome.,function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity.,similarity:Belongs to the KIF1-binding protein family.,subunit:Interacts with KIF1B.,tissue specificity:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined, as well as in	Cell Pathway	Cytoplasm, cytoskeleton .
syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome.,function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity.,similarity:Belongs to the KIF1-binding protein family.,subunit:Interacts with KIF1B.,tissue specificity:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in	Tissue Specificity	regions examined. Moderate expressed at intermediate level in all other adult tissues examined, as well as in fetal liver and brain. Not expressed in blood
	Function	syndrome (GOSHS) [MIM:609460]. GOSHS is characterized by microcephaly, mental retardation and facial dysmorphism, as well as phenotypes related to Hirschsprung disease syndrome.,function:Required for organization of axonal microtubules, and axonal outgrowth and maintenance during peripheral and central nervous system development. Regulates mitochondrial transport by modulating KIF1B motor activity.,similarity:Belongs to the KIF1-binding protein family.,subunit:Interacts with KIF1B.,tissue specificity:Highly expressed in heart, brain, ovary, testis, spinal cord and all specific brain regions examined. Moderate

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






Background	This gene encodes a kinesin family member 1 binding protein that is characterized by two tetratrico peptide repeats. The encoded protein localizes to the mitochondria and may be involved in regulating transport of the mitochondria. Mutations in this gene are associated with Goldberg-Shprintzen megacolon syndrome. [provided by RefSeq, Mar 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

fetal liver and brain. Not expressed in blood leukocytes.

## **Products Images**



Western Blot analysis of various cells using KBP mouse mAb

Nanjing BYabscience technology Co.,Ltd