



## AFAF mouse mAb

Catalog No   BYmab-11800     Isotype   IgG     Reactivity   Human;Rat;Mouse;     Applications   WB     Gene Name   AFAF C9orf11     Protein Name   AFAF     Immunogen   Synthesized peptide derived from human AFAF AA range; 223-273     Specificity   This antibody detects endogenous levels of AFAF at Human     Formulation   Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.     Source   Monoclonal, Mouse, IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Plution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   -     Cell Pathway   Cytoplasmic vesicle, secretory vesicle, secreto		
Reactivity Human;Rat;Mouse;   Applications WB   Gene Name AFAF C9orf11   Protein Name AFAF   Immunogen Synthesized peptide derived from human AFAF AA range: 223-273   Specificity This antibody detects endogenous levels of AFAF at Human   Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type 1 membrane protein . Cytoplasmic vesicle, acrosome nembrane ; Single-pass type 1 membrane protein . Cytoplasmic vesicle, acrosome nembrane ; Single-pass type 1 membrane protein . In the anterior acrosome are gion enriched on but the inner and outer acrosomal membrane but minimal on the outer acrosomal membrane ; Single-pass type 1 membrane protein . In the anterior acrosome are in contrast in the posterior . In the anterior acrosome eigen on the inner and outer acrosomal membrane but minimal on the outer acrosomal membrane protein . I	Catalog No	BYmab-11800
ApplicationsWBGene NameAFAF C9orf11Protein NameAFAFImmunogenSynthesized peptide derived from human AFAF AA range: 223-273SpecificityThis antibody detects endogenous levels of AFAF at HumanFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse, IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCytoplasmic vesicle, secretory vesicle, acrosome membrane : Single-pass type 1 membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane : Single-pass type 1 membrane protein . Cytoplasmic vesicle, secretory vesicle	Isotype	lgG
Gene Name   AFAF C9orf11     Protein Name   AFAF     Immunogen   Synthesized peptide derived from human AFAF AA range: 223-273     Specificity   This antibody detects endogenous levels of AFAF at Human     Formulation   Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.     Source   Monoclonal, Mouse,IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane in contrast in the posterior acrosome inner membrane is on the outer acrosomal membrane stretor acrosome inner membrane is in on the outer acrosomal membrane is norther acrosomal membranes.     Tissue Specificity   Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.,	Reactivity	Human;Rat;Mouse;
Protein Name   AFAF     Immunogen   Synthesized peptide derived from human AFAF AA range: 223-273     Specificity   This antibody detects endogenous levels of AFAF at Human     Formulation   Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.     Source   Monoclonal, Mouse,IgG     Purification   The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.     Dilution   WB 1:500-2000     Concentration   1 mg/ml     Purity   ≥90%     Storage Stability   -20°C/1 year     Synonyms   Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane in contrast in the posterior acrosome region, enriched on the inner acrosomal membranes.     Tissue Specificity   Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.     Function   function.Involved in acrosome biogenesis.tissue specificity.Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Applications	WB
ImmunogenSynthesized peptide derived from human AFAF AA range: 223-273SpecificityThis antibody detects endogenous levels of AFAF at HumanFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourceMonoclonal, Mouse,IgGPurificationThe antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearCell PathwayCytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, secretory vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, secretory vesicle, acrosome membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome membrane protein . In the anterior acrosome outer membrane; in contras in the posterior acrosome region enriched on both the inner and outer acrosomal membrane; minal on the outer acrosomal membrane; in contras of and blood.Functionfunction.Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Gene Name	AFAF C9orf11
Specificity This antibody detects endogenous levels of AFAF at Human   Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type 1 membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome region enriched on both the inner and outer acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Protein Name	AFAF
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   Source Monoclonal, Mouse,IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome membrane protein . In the anterior acrosome or ergion, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane s.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.,   Function function:Involved in acrosome biogenesis, tissue specificity:Isoform 1 highly expressed at low levels in skin and blood.,	Immunogen	Synthesized peptide derived from human AFAF AA range: 223-273
Source Monoclonal, Mouse, IgG   Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . In the arterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane protein . In the arterior acrosome region enriched on both the inner and outer acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis., tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Specificity	This antibody detects endogenous levels of AFAF at Human
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cotyplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome region enriched on both the inner and outer acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis.tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.   Dilution WB 1:500-2000   Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . In the anterior acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis, tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml   Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I   Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . In the anterior acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Purification	
Purity ≥90%   Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band -20°C/1 year   Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome region enriched on both the inner and outer acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis.tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.,	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year   Synonyms -20°C/1 year   Observed Band Cell Pathway   Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome membrane; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed at low levels in skin and blood.,	Concentration	1 mg/ml
Synonyms   Observed Band   Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome region enriched on both the inner and outer acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Purity	≥90%
Observed Band   Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome region enriched on both the inner and outer acrosomal membranes.   Tissue Specificity Isoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.   Function function:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Storage Stability	-20°C/1 year
Cell PathwayCytoplasmic vesicle, secretory vesicle, acrosome membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome region enriched on both the inner and outer acrosomal membranes.Tissue SpecificityIsoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.Functionfunction:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.	Synonyms	
membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome region enriched on both the inner and outer acrosomal membranesTissue SpecificityIsoform 1 is highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.Functionfunction:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.,	Observed Band	
Function function:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.,	Cell Pathway	membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome inner membrane ; Single-pass type I membrane protein . Cytoplasmic vesicle, secretory vesicle, acrosome outer membrane ; Single-pass type I membrane protein . In the anterior acrosome region, enriched on the inner acrosomal membrane but minimal on the outer acrosomal membrane; in contrast in the posterior acrosome
expressed in testis. Isoform 2 is expressed at low levels in skin and blood.,	Tissue Specificity	
Background	Function	function:Involved in acrosome biogenesis.,tissue specificity:Isoform 1 highly expressed in testis. Isoform 2 is expressed at low levels in skin and blood.,
	Background	

Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998



attention


matters needing



## Avoid repeated freezing and thawing!

**Usage suggestions** This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Nanjing BYabscience technology Co.,Ltd