



## ERLN1 mouse mAb

Catalog No	BYmab-12104
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	ERLIN1 C10orf69 KE04 KEO4 SPFH1
Protein Name	ERLN1
Immunogen	Synthesized peptide derived from human ERLN1 AA range: 185-235
Specificity	This antibody detects endogenous levels of ERLN1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
	The antibody was affinity-purified from mouse antiserum by
Purification	affinity-chromatography using epitope-specific immunogen.
Purification  Dilution	
	affinity-chromatography using epitope-specific immunogen.
Dilution	affinity-chromatography using epitope-specific immunogen. WB 1:500-2000
Dilution  Concentration	affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000  1 mg/ml
Dilution  Concentration  Purity	affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000  1 mg/ml  ≥90%
Dilution  Concentration  Purity  Storage Stability	affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000  1 mg/ml  ≥90%
Dilution  Concentration  Purity  Storage Stability  Synonyms	affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000  1 mg/ml  ≥90%
Dilution  Concentration  Purity  Storage Stability  Synonyms  Observed Band	affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000  1 mg/ml ≥90% -20°C/1 year
Dilution Concentration Purity Storage Stability Synonyms Observed Band Cell Pathway	affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000  1 mg/ml  ≥90%  -20°C/1 year  Endoplasmic reticulum membrane ; Single-pass type II membrane protein . Associated with lipid raft-like domains of the endoplasmic reticulum membrane.  Expressed in heart, placenta, liver, kidney, pancreas, prostate, testis, ovary and
Dilution  Concentration  Purity  Storage Stability  Synonyms  Observed Band  Cell Pathway  Tissue Specificity	affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000  1 mg/ml  ≥90%  -20°C/1 year  Endoplasmic reticulum membrane ; Single-pass type II membrane protein . Associated with lipid raft-like domains of the endoplasmic reticulum membrane.  Expressed in heart, placenta, liver, kidney, pancreas, prostate, testis, ovary and small intestine.  similarity:Belongs to the band 7/mec-2 family.,subcellular location:Associated with lipid raft-like domains of the endoplasmic reticulum membrane.,tissue specificity:Expressed in heart, placenta, liver, kidney, pancreas, prostate, testis,

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western Blot analysis of various cells using ERLN1 mouse mAb