



PUR6 mouse mAb

Catalog No	BYmab-12117
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	PAICS ADE2 AIRC PAIS
Protein Name	PUR6
Immunogen	Synthesized peptide derived from human PUR6 AA range: 107-157
Specificity	This antibody detects endogenous levels of PUR6 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	cytoplasm,cytosol,cell-cell adherens junction,membrane,extracellular exosome,
Tissue Specificity	
Function	catalytic activity:5-amino-1-(5-phospho-D-ribose)imidazole-4-carboxylate = 5-amino-1-(5-phospho-D-ribose)imidazole + CO(2).,catalytic activity:ATP + 5-amino-1-(5-phospho-D-ribose)imidazole-4-carboxylate + L-aspartate = ADP + phosphate + (S)-2-(5-amino-1-(5-phospho-D-ribose)imidazole-4-carboxamido)succinate.,path way:Purine metabolism; IMP biosynthesis via de novo pathway; 5-amino-1-(5-phospho-D-ribose)imidazole-4-carboxamide from N(2)-formyl-N(1)-(5-phospho-D-ribose)glycinamide: step 3/5.,pathway:Purine metabolism; IMP biosynthesis via de novo pathway; 5-amino-1-(5-phospho-D-ribose)imidazole-4-carboxamide from N(2)-formyl-N(1)-(5-phospho-D-ribose)glycinamide: step 4/5.,similarity:In the C-terminal section; belongs to the AIR carboxylase family.,similarity:In the

Nanjing BYabscience technology Co.,Ltd



N-terminal section; belongs to the SAICAR synthetase family.,subunit:Homooctamer.,

Background

This gene encodes a bifunctional enzyme containing phosphoribosylaminoimidazole carboxylase activity in its N-terminal region and phosphoribosylaminoimidazole succinocarboxamide synthetase in its C-terminal region. It catalyzes steps 6 and 7 of purine biosynthesis. The gene is closely linked and divergently transcribed with a locus that encodes an enzyme in the same pathway, and transcription of the two genes is coordinately regulated. The human genome contains several pseudogenes of this gene. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using PUR6 mouse mAb