



HSF2B mouse mAb

Catalog No BYmab-12186 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name HSF2BP Protein Name HSF2B Immunogen Synthesized peptide derived from human HSF2B AA range: 201-251 Specificity This antibody detects endogenous levels of HSF2B At Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function. May be involved in modulating HSF2 activation in testis. PTM:3umyolyated by UBE21 in response to MEKK1-mediated stmuli, subunit/Associates with HSF2. The interaction seems to occur between the HSF2BP, tissue specificity Testis specific. <		
Reactivity Human;Rat;Mouse; Applications WB Gene Name HSF2BP Protein Name HSF2B Immunogen Synthesized peptide derived from human HSF2B AA range: 201-251 Specificity This antibody detects endogenous levels of HSF2B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis., PTM. Sumoylated by UBE21 in response to MEKK1-mediated stimuli. subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity. Testis specific. Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Catalog No	BYmab-12186
Applications Gene Name HSF2BP Protein Name HSF2B Immunogen Synthesized peptide derived from human HSF2B AA range: 201-251 Specificity This antibody detects endogenous levels of HSF2B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis., PTM:Sumoylated by UBE2! in response to MEKK1-mediated stimuli., subunit:Associates with HSF2. The interaction seems to occur between the timerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity. Testis specific. Background	Isotype	IgG
Gene Name HSF2BP Protein Name HSF2B Immunogen Synthesized peptide derived from human HSF2B AA range: 201-251 Specificity This antibody detects endogenous levels of HSF2B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm. Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis., PTM:Sumoylated by UBE21 in response to MEKK1-mediated stimuli, subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-Herminal hydrophilic region of HSF2BP may therefore be involved in modulating HSF2 ascoviates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the N-Herminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [Provided by RefSeq, Jul	Reactivity	Human;Rat;Mouse;
Protein Name HSF2B Immunogen Synthesized peptide derived from human HSF2B AA range: 201-251 Specificity This antibody detects endogenous levels of HSF2B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2! in response to MEKK1-mediated stimuli, subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity: Testis specific. Background before the trimerization domain of HSF2 and the A-terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP may therefore be involved in modulating HSF2 activation (HSF2BP) activation (HSF3BP) may therefore be involved in modulating HSF2 activation (HSF3BP) may therefore be involved in modulating HSF2 activation (HSF3BP) may	Applications	WB
Immunogen Synthesized peptide derived from human HSF2B AA range: 201-251 Specificity This antibody detects endogenous levels of HSF2B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity:Testis specific. Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the Amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP MSY therefore be involved in modulating HSF2 activation, Glycologi	Gene Name	HSF2BP
Specificity This antibody detects endogenous levels of HSF2B at Human Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis. PTM:Sumoylated by UBE21 in response to MEKK1-mediated stimuli, subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RFS2BP may therefore be involved in mediating PSF2 activation. [provided by RFS2BP may therefore be involved in modulating HSF2 activation. [provided by RFS2BP may therefore be involved in modulating HSF2 activation. [provided by RFS2BP may therefore be involved in modulating HSF2 activation. [provided by RFS2BP, Juli	Protein Name	HSF2B
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis. ,PTM:Sumoylated by UBE21 in response to MEKK1-mediated stimulii, subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2BP and the N-terminal hydrophilic region of HSF2BP, that comprises two leucine zipper motifis. HSF2BP may therefore be involved in modulating HSF2 and the amino terminal hydrophilic region of HSF2BP hay therefore be involved in modulating HSF2 and the amino terminal hydrophilic region of HSF2BP into comprises two leucine zipper motifis. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Immunogen	Synthesized peptide derived from human HSF2B AA range: 201-251
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2l in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity. Testis specific. Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Specificity	This antibody detects endogenous levels of HSF2B at Human
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis., PTM:Sumoylated by UBE2! in response to MEKK1-mediated stimuli., subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity:Testis specific. Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jull	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the frimerization domain of HSF2 and the Amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes. Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis., PTM:Sumoylated by UBE21 in response to MEKK1-mediated stimuli., subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP., tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP.,tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP, tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP.,tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Synonyms	
Tissue Specificity Testis specific. Overexpressed in some tumors (PubMed:31242413). Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP.,tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Observed Band	
Function function:May be involved in modulating HSF2 activation in testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP.,tissue specificity:Testis specific., HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Cell Pathway	Cytoplasm . Chromosome . Localizes on double-strand breaks (DSBs) in mitotic and meiotic chromosomes
testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of HSF2BP.,tissue specificity:Testis specific., Background HSF2 binding protein (HSF2BP) associates with HSF2. The interaction occurs between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Tissue Specificity	Testis specific. Overexpressed in some tumors (PubMed:31242413).
between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may therefore be involved in modulating HSF2 activation. [provided by RefSeq, Jul	Function	testis.,PTM:Sumoylated by UBE2I in response to MEKK1-mediated stimuli.,subunit:Associates with HSF2. The interaction seems to occur between the trimerization domain of HSF2 and the N-terminal hydrophilic region of
	Background	between the trimerization domain of HSF2 and the amino terminal hydrophilic region of HSF2BP that comprises two leucine zipper motifs. HSF2BP may

Nanjing BYabscience technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

