



KNTC1 mouse mAb

Catalog No	BYmab-12264
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	KNTC1 KIAA0166
Protein Name	KNTC1
Immunogen	Synthesized peptide derived from human KNTC1 AA range: 1343-1393
Specificity	This antibody detects endogenous levels of KNTC1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm. Nucleus. Chromosome, centromere, kinetochore. Cytoplasm, cytoskeleton, spindle. Dynamic pattern of localization during the cell cycle. At interphase, uniformly distributed throughout the cytoplasm and nucleus. By prophase and until late stages of prometaphase, a fraction of the total pool is concentrated at kinetochores. By metaphase, detected at kinetochores, along spindle fibers and most prominently at the poles. By late anaphase until the end of telophase, no longer detectable on kinetochores or along spindle fibers, but still present at the spindle poles.
Tissue Specificity	High expression in testis.
Function	function:Essential component of the mitotic checkpoint, which prevents cells from prematurely exiting mitosis. Required for the assembly of the dynein-dynactin and MAD1-MAD2 complexes onto kinetochores.,subcellular location:Dynamic pattern of localization during the cell cycle. At interphase, uniformly distributed throughout the cytoplasm and nucleus. By prophase and until late stages of prometaphase, a fraction of the total pool is concentrated at kinetochores. By metaphase, detected

Nanjing BYabscience technology Co.,Ltd



at kinetochores, along spindle fibers and most prominently at the poles. By late anaphase until the end of telophase, no longer detectable on kinetochores or along spindle fibers, but still present at the spindle poles.,subunit:Interacts with ZW10. This interaction is required for stable association with the kinetochore. Component of the RZZ complex composed of KNTC1/ROD, ZW10 and ZWILCH.,tissue specificity

Background

This gene encodes a protein that is one of many involved in mechanisms to ensure proper chromosome segregation during cell division. Experimental evidence indicated that the encoded protein functioned in a similar manner to that of the Drosophila rough deal protein. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using KNTC1 mouse mAb