

## PCDBC mouse mAb

Catalog No         BYmab-12275           Isotype         IgG           Reactivity         Human:Rat;Mouse;           Applications         WB           Gene Name         PCDHB12           Protein Name         PCDBC           Immunogen         Synthesized peptide derived from human PCDBC AA range: 555-605           Specificity         This antibody detects endogenous levels of PCDBC at Human           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Cell Pathway         Cell membrane; Single-pass type I membrane protein .           Tissue Specificity         Function           Function         function: Potential calcium-dependent cell-adhesion protein May be involved in the establishment and maintenance of specific neuronal connections in the brain , similarity: Contains 8 cadherin domains.           Background         This gene is a member of the protocadherin beta gene cluster, one of		
Reactivity Human;Rat;Mouse;  Applications WB  Gene Name PCDHB12  Protein Name PCDBC  Immunogen Synthesized peptide derived from human PCDBC AA range: 555-605  Specificity This antibody detects endogenous levels of PCDBC at Human  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse,IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain, similarity:Contains 6 cadherin domains.,  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters tandemly linked on chromosome five. The gene clusters clemonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters from others in the cadherin superfamily. The extracellular comains interact in a homosphilic manner to specify differential cell-connections. Unlike interact in a homosphilic manner to specify differential cell-connections. Unlike interact in a homosphilic manner to specify differential cell-connections. Unlike interact in a homosphilic manner to specify differential cell-connections. Unlike	Catalog No	BYmab-12275
Applications  Gene Name PCDHB12 Protein Name PCDBC  Immunogen Synthesized peptide derived from human PCDBC AA range: 555-605  Specificity This antibody detects endogenous levels of PCDBC at Human  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain., similarity:Contains 6 cadherin domains.,  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta clusters contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular concentions. Unlike	Isotype	IgG
Gene Name         PCDHB12           Protein Name         PCDBC           Immunogen         Synthesized peptide derived from human PCDBC AA range: 555-605           Specificity         This antibody detects endogenous levels of PCDBC at Human           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, lgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Cell Pathway         Cell membrane; Single-pass type I membrane protein.           Tissue Specificity         Function           Function         function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain. similarity. Contains 6 cadherin domains.           Background         This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, aca hencoding 6 catherin domains and a cytoplasmic	Reactivity	Human;Rat;Mouse;
Protein Name PCDBC  Immunogen Synthesized peptide derived from human PCDBC AA range: 555-605  Specificity This antibody detects endogenous levels of PCDBC at Human  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane ; Single-pass type I membrane protein .  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain., similarity:Contains 6 cadherin domains.  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell connections. Unlike interact in a homophilic manner to specify differential cell connections. Unlike	Applications	WB
Immunogen         Synthesized peptide derived from human PCDBC AA range: 555-605           Specificity         This antibody detects endogenous levels of PCDBC at Human           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Monoclonal, Mouse, IgG           Purification         The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Cell Pathway         Cell membrane; Single-pass type I membrane protein.           Tissue Specificity         Function           Function         function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,           Background         This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify	Gene Name	PCDHB12
Specificity This antibody detects endogenous levels of PCDBC at Human  Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact from others in the cadherin superfamily. The extracellular cometions. Unlike	Protein Name	PCDBC
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Monoclonal, Mouse, IgG  Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane ; Single-pass type I membrane protein .  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Immunogen	Synthesized peptide derived from human PCDBC AA range: 555-605
Source       Monoclonal, Mouse,IgG         Purification       The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Cell Pathway         Cell Pathway       Cell membrane ; Single-pass type I membrane protein .         Tissue Specificity       Function         function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain, similarity:Contains 6 cadherin domains.         Background       This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Specificity	This antibody detects endogenous levels of PCDBC at Human
Purification  The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  Cell Pathway  Cell membrane ; Single-pass type I membrane protein .  Tissue Specificity  Function  function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane ; Single-pass type I membrane protein .  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Purification	•
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Dilution	WB 1:500-2000
Synonyms  Observed Band  Cell Pathway  Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function  function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Concentration	1 mg/ml
Synonyms  Observed Band  Cell Pathway  Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function  function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Purity	≥90%
Cell Pathway  Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function  function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Storage Stability	-20°C/1 year
Cell Pathway  Cell membrane; Single-pass type I membrane protein.  Tissue Specificity  Function  function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  Background  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Synonyms	
Tissue Specificity  Function  function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Observed Band	
Function  function:Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Cell Pathway	Cell membrane ; Single-pass type I membrane protein .
the establishment and maintenance of specific neuronal connections in the brain.,similarity:Contains 6 cadherin domains.,  This gene is a member of the protocadherin beta gene cluster, one of three related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Tissue Specificity	
related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike	Function	the establishment and maintenance of specific neuronal connections in the
	Background	related gene clusters tandemly linked on chromosome five. The gene clusters demonstrate an unusual genomic organization similar to that of B-cell and T-cell receptor gene clusters. The beta cluster contains 16 genes and 3 pseudogenes, each encoding 6 extracellular cadherin domains and a cytoplasmic tail that deviates from others in the cadherin superfamily. The extracellular domains interact in a homophilic manner to specify differential cell-cell connections. Unlike

Nanjing BYabscience technology Co.,Ltd


网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





	only one large exon, not sharing common 3' exons as expected. These neural cadherin-like cell adhesion proteins are integral plasma membrane proteins. Their specific functions are unknown but they most likely play a critical role in the establishment and function of specific cell-cell neural connections. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658