



# PRGC2 mouse mAb

Catalog No	BYmab-12361
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	PPARGC1B PERC PGC1 PGC1B PPARGC1
Protein Name	PRGC2
Immunogen	Synthesized peptide derived from human PRGC2 AA range: 829-879
Specificity	This antibody detects endogenous levels of PRGC2 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	113kD
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitous with higher expression in heart, brain and skeletal muscle.
Function	domain:Contains 2 Leu-Xaa-Xaa-Leu-Leu (LXXLL) motif, which are usually required for the association with nuclear receptors.,function:Plays a role of stimulator of transcription factors and nuclear receptors activities. Activates transcripnal activity of estrogen receptor alpha, nuclear respiratory factor 1 (NRF1) and glucocorticoid receptor in the presence of glucocorticoids. May play a role in constitutive non-adrenergic-mediated mitochondrial biogenesis as suggested by increased basal oxygen consumption and mitochondrial number when overexpressed. May be involved in fat oxidation and non-oxidative glucose metabolism and in the regulation of energy expenditure.,induction:Repressed by saturated fatty acids such as palmitate and stearate in skeletal muscle cells. Induced by insulin and reduced by aging in skeletal muscle biopsies. Down-regulated in type 2 diabetes mellitus subjects as w

Nanjing BYabscience technology Co.,Ltd



<b>Background</b>	The protein encoded by this gene stimulates the activity of several transcription factors and nuclear receptors, including estrogen receptor alpha, nuclear respiratory factor 1, and glucocorticoid receptor. The encoded protein may be involved in fat oxidation, non-oxidative glucose metabolism, and the regulation of energy expenditure. This protein is downregulated in prediabetic and type 2 diabetes mellitus patients. Certain allelic variations in this gene increase the risk of the development of obesity. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2010],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

