



# LC3 A/B mouse mAb

Catalog No	BYmab-12495
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	LC3 A/B
Protein Name	LC3 A/B
Immunogen	Synthesized peptide derived from human LC3 A/B
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat LC3 A/B
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	14,16kD
Cell Pathway	Cytoplasmic vesicle, autophagosome membrane ; Lipid-anchor . Endomembrane system ; Lipid-anchor . Cytoplasm, cytoskeleton . LC3-II binds to the autophagic membranes. .
Tissue Specificity	Most abundant in heart, brain, liver, skeletal muscle and testis but absent in thymus and peripheral blood leukocytes.
Function	autophagic vacuole formation, proteolysis, autophagy, vacuole organization, macromolecule catabolic process, cellular response to starvation, response to extracellular stimulus, macroautophagy, modification-dependent protein catabolic process, protein catabolic process, response to nutrient levels, cellular response to extracellular stimulus, cellular response to nutrient levels, cellular response to stress, response to starvation, modification-dependent macromolecule catabolic process, cellular protein catabolic process, cellular macromolecule catabolic process,proteolysis involved in cellular protein catabolic process,

Nanjing BYabscience technology Co.,Ltd



#### Background

function:Probably involved in formation of autophagosomal vacuoles (autophagosomes).,PTM:The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II.,similarity:Belongs to the MAP1 LC3 family.,subcellular location:LC3-II binds to the autophagic membranes.,subunit:3 different light chains, LC1, LC2 and LC3, can associate with MAP1A and MAP1B proteins.,tissue specificity:Most abundant in heart, brain, liver, skeletal muscle and testis but absent in thymus and peripheral blood leukocytes.,

#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images

Nanjing BYabscience technology Co.,Ltd

网址: [www.njbybio.com](http://www.njbybio.com)

官方热线: 025-5229-8998

监督电话: 15950492658