




TIFA mouse mAb

Catalog No BYmab-12531 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name TIFA T2BP Protein Name TIFA Immunogen Synthesized peptide derived from human TIFA Specificity This antibody detects endogenous levels of TIFA at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein) PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation		
Reactivity Applications WB Gene Name TIFA T2BP Protein Name ITIFA Immunogen Synthesized peptide derived from human TIFA Specificity This antibody detects endogenous levels of TIFA at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Catalog No	BYmab-12531
Applications Gene Name TIFA T2BP Protein Name TIFA Immunogen Synthesized peptide derived from human TIFA Specificity This antibody detects endogenous levels of TIFA at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm. Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation. Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name Protein Name TIFA T2BP Protein Name TIFA Immunogen Synthesized peptide derived from human TIFA Specificity This antibody detects endogenous levels of TIFA at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with Iysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Reactivity	Human; Mouse;Rat
Protein Name Immunogen Synthesized peptide derived from human TIFA Specificity This antibody detects endogenous levels of TIFA at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Applications	WB
Immunogen Synthesized peptide derived from human TIFA Specificity This antibody detects endogenous levels of TIFA at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation. Tissue Specificity Function Background Matters needing Avoid repeated freezing and thawing!	Gene Name	TIFA T2BP
Specificity This antibody detects endogenous levels of TIFA at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Protein Name	TIFA
Formulation Liquid in PBS containing 50% glycerol, and 0.13% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human TIFA
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with Iysosomal marker LAMP2 following homooligomerization and subsequent activation. Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of TIFA at Human, Mouse,Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.13% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation . Tissue Specificity Function Background Avoid repeated freezing and thawing!	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms TRAF-interacting protein with FHA domain-containing protein A (Putative MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Matters needing Avoid repeated freezing and thawing!	Purity	≥90%
MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20) (TRAF2-binding protein) Observed Band Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation Tissue Specificity Function Background Matters needing Avoid repeated freezing and thawing!	Synonyms	MAPK-activating protein PM14) (Putative NF-kappa-B-activating protein 20)
homooligomerization and subsequent activation Tissue Specificity Function Background matters needing Avoid repeated freezing and thawing!	Observed Band	
Function Background matters needing Avoid repeated freezing and thawing!	Cell Pathway	Cytoplasm . Colocalizes with lysosomal marker LAMP2 following homooligomerization and subsequent activation
Background matters needing Avoid repeated freezing and thawing!	Tissue Specificity	
matters needing Avoid repeated freezing and thawing!	Function	
matters needing Avoid repeated freezing and thawing! attention	Background	
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd








Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using TIFA mouse mAb

Nanjing BYabscience technology Co.,Ltd