






Catalog No	BYmab-12551
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	TPPP TPPP1
Protein Name	TPPP
Immunogen	Synthesized peptide derived from human TPPP
Specificity	This antibody detects endogenous levels of TPPP at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.39% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Tubulin polymerization-promoting protein (TPPP) (25 kDa brain-specific protein) (TPPP/p25) (p24) (p25-alpha)
Observed Band	
Cell Pathway	Golgi outpost . Cytoplasm, cytoskeleton, microtubule organizing center . Cytoplasm, cytoskeleton . Nucleus . Cytoplasm, cytoskeleton, spindle . Specifically localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, which shapes dendrite morphology by functioning as sites of acentrosomal microtubule nucleation (By similarity). Mainly localizes to the cytoskeleton (PubMed:18028908). Also found in the nucleus; however, nuclear localization is unclear and requires additional evidences (PubMed:18028908). Localizes to glial Lewy bodies in the brains of individuals with synucleinopathies (PubMed:15590652, PubMed:17027006). During mitosis, colocalizes with LIMK2 at the mitotic spindle (PubMed:22328514).
Tissue Specificity	Widely expressed.
Function	function:Promotes in vitro the polymerization of tubulin into double-walled tubules and polymorphic aggregates or bundled stabilized microtubules blocks. When
	overexpressed, inhibits mitotic spindle assembly and nuclear envelope

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








	breakdown, apparently without affecting other cellular events.,PTM:Phosphorylated by TPK II. Poor substrate for GSK3.,similarity:Belongs to the TPPP family.,subunit:May form dimers. Binds tubulin; binding is inhibited by GTP. Interacts with GSK3.,tissue specificity:Widely expressed.,
Background	
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658