



PLEKHG5 mouse mAb

Catalog No BYmab-12573 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name PLEKHG5 KIAA0720 Protein Name PLEKHG5 Immunogen Synthesized peptide derived from human PLEKHG5 Specificity This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cytoplasm. Cytoplasm, perinuclear region. Cell membrane. Cell junction Cell projection, lamellipodium. Predominantity cytoplasmic, however when endothelial cells and returnated with lysophosphalidic acid, PLEKHG5 is forwered when endothelial cells, and relocalizes to cytoplasmic vesicle and the leading endothelial cells, and relocalizes to cytoplasmic vesicle and the leading endothelial cells, an		
Reactivity Human; Mouse;Rat Applications WB Gene Name PLEKHG5 KIAA0720 Protein Name PLEKHG5 Immunogen Synthesized peptide derived from human PLEKHG5 Specificity This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vessicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Catalog No	BYmab-12573
Applications WB Gene Name PLEKHG5 KIAA0720 Protein Name PLEKHG5 Immunogen Synthesized peptide derived from human PLEKHG5 Specificity This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm, Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Isotype	IgG
Gene Name PLEKHG5 KIAA0720 Protein Name PLEKHG5 Immunogen Synthesized peptide derived from human PLEKHG5 Specificity This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cytoplasm . Cytoplasm , perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present i	Reactivity	Human; Mouse;Rat
Protein Name PLEKHG5 Immunogen Synthesized peptide derived from human PLEKHG5 Specificity This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cytoplasm . Cytoplasm , perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell-junctions in quiessent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in gliobl	Applications	WB
Immunogen Synthesized peptide derived from human PLEKHG5 Specificity This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cytoplasm . Cytoplasm , perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphaticia acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Gene Name	PLEKHG5 KIAA0720
Specificity This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm , perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantily cytoplasmic, however when endothelial cells are stimulated with tysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells . Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Protein Name	PLEKHG5
Formulation Liquid in PBS containing 50% glycerol, and 0.64% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphaticic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Immunogen	Synthesized peptide derived from human PLEKHG5
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm Cytoplasm, perinuclear region Cell membrane Cell junction Cell projection, lamellipodium Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Specificity	This antibody detects endogenous levels of PLEKHG5 at Human, Mouse,Rat
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Formulation	Liquid in PBS containing 50% glycerol, and 0.64% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane . Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells . Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Concentration	1 mg/ml
Pleckstrin homology domain-containing family G member 5 (PH domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Purity	≥90%
domain-containing family G member 5) (Guanine nucleotide exchange factor 720) (GEF720) Observed Band Cell Pathway Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Storage Stability	-20°C/1 year
Cytoplasm . Cytoplasm, perinuclear region . Cell membrane . Cell junction . Cell projection, lamellipodium . Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Synonyms	domain-containing family G member 5) (Guanine nucleotide exchange factor 720)
projection, lamellipodium. Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of lamellipodia in migrating endothelial cells. Tissue Specificity Predominantly expressed in the peripheral nervous system and brain. Highest expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Observed Band	
expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed in glioblastoma (GBM) cell lines.	Cell Pathway	projection, lamellipodium. Predominantly cytoplasmic, however when endothelial cells are stimulated with lysophosphatidic acid, PLEKHG5 is found in perinuclear regions and at the cell membrane. Localizes at cell-cell junctions in quiescent endothelial cells, and relocalizes to cytoplasmic vesicle and the leading edge of
Function	Tissue Specificity	expression is observed in heart, lung, kidney, testis and moderate expression is present in spleen, pancreas, skeletal muscle, ovary and liver. Weakly expressed
	Function	

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images