



# CSP Monoclonal Antibody

<b>Catalog No</b>	BYmab-12706
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	DNAJC5
<b>Protein Name</b>	DnaJ homolog subfamily C member 5
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human DNAJC5. AA range:131-180
<b>Specificity</b>	CSP Monoclonal Antibody detects endogenous levels of CSP protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	DNAJC5; CSP; DnaJ homolog subfamily C member 5; Cysteine string protein; CSP
<b>Observed Band</b>	30kD
<b>Cell Pathway</b>	Cytoplasm, cytosol . Membrane ; Lipid-anchor . Cytoplasmic vesicle, secretory vesicle, chromaffin granule membrane . Melanosome . Cell membrane . The association with membranes is regulated by palmitoylation (By similarity). Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). .
<b>Tissue Specificity</b>	Expressed in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain and heart.
<b>Function</b>	function:May have an important role in presynaptic function. May be involved in calcium-dependent neurotransmitter release at nerve endings.,PTM:Fatty acylated. Heavily palmitoylated in the cysteine string motif.,similarity:Contains 1 J domain.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,tissue specificity:Expressed in pancreas, kidney, skeletal muscle, liver, lung, placenta, brain and heart.,

**Nanjing BYabscience technology Co.,Ltd**



## Background

This gene is a member of the J protein family. J proteins function in many cellular processes by regulating the ATPase activity of 70 kDa heat shock proteins. The encoded protein plays a role in membrane trafficking and protein folding, and has been shown to have anti-neurodegenerative properties. The encoded protein is known to play a role in cystic fibrosis and Huntington's disease. A pseudogene of this gene is located on the short arm of chromosome 8. [provided by RefSeq, Nov 2010],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using CSP Monoclonal Antibody