




GAP-43 Monoclonal Antibody

Catalog No	BYmab-12727
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GAP43
Protein Name	Neuromodulin
Immunogen	The antiserum was produced against synthesized peptide derived from human GAP43. AA range:8-57
Specificity	GAP-43 Monoclonal Antibody detects endogenous levels of GAP-43 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GAP43; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46
Observed Band	43kD
Cell Pathway	Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, growth cone membrane; Peripheral membrane protein; Cytoplasmic side. Cell junction, synapse. Cell projection, filopodium membrane; Peripheral membrane protein. Perikaryon. Cell projection, dendrite. Cell projection, axon. Cytoplasm. Cytoplasmic surface of growth cone and synaptic plasma membranes.
Tissue Specificity	Alzheimer cortex,Brain,Subthalamic nucleus,
Function	function: This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons., online information: Gap-43 entry, PTM: Phosphorylation of this protein by a protein kinase C is specifically correlated with certain forms of synaptic plasticity., similarity: Belongs to the neuromodulin family., similarity: Contains 1 IQ

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








	plasma membranes.,subunit:Binds calmodulin with a greater affinity in the absence of Ca(2+) than in its presence.,
Background	The protein encoded by this gene has been termed a 'growth' or 'plasticity' protein because it is expressed at high levels in neuronal growth cones during development and axonal regeneration. This protein is considered a crucial component of an effective regenerative response in the nervous system. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using GAP-43 Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd