



# Netrin-1 Monoclonal Antibody

<b>Catalog No</b>	BYmab-12749
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat;Monkey
<b>Applications</b>	WB
<b>Gene Name</b>	NTN1
<b>Protein Name</b>	Netrin-1
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human Netrin-1. AA range:484-533
<b>Specificity</b>	Netrin-1 Monoclonal Antibody detects endogenous levels of Netrin-1 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	NTN1; NTN1L; Netrin-1; Epididymis tissue protein Li 131P
<b>Observed Band</b>	75kD
<b>Cell Pathway</b>	Secreted . Cytoplasm . Mainly secreted. .
<b>Tissue Specificity</b>	Widely expressed in normal adult tissues with highest levels in heart, small intestine, colon, liver and prostate. Reduced expression in brain tumors and neuroblastomas. Expressed in epididymis (at protein level).
<b>Function</b>	disease:Defects in NTN1 are associated with some forms of neuroblastomas.,function:Netrins control guidance of CNS commissural axons and peripheral motor axons. Its association with either DCC or some UNC5 receptors will lead to axon attraction or repulsion, respectively. It also serve as a survival factor via its association with its receptors which prevent the initiation of apoptosis. Involved in colorectal tumorigenesis by regulating apoptosis.,similarity:Contains 1 laminin N-terminal domain.,similarity:Contains 1 NTR domain.,similarity:Contains 3 laminin EGF-like domains.,subunit:Binds to its receptors; DCC, UNC5A, UNC5B, UNC5C and probably UNC5D.,tissue specificity:Widely expressed in normal adult tissues with highest levels in heart,

**Nanjing BYabscience technology Co.,Ltd**



small intestine, colon, liver and prostate. Reduced expression in brain tumors and neuroblastomas.,

## Background

Netrin is included in a family of laminin-related secreted proteins. The function of this gene has not yet been defined; however, netrin is thought to be involved in axon guidance and cell migration during development. Mutations and loss of expression of netrin suggest that variation in netrin may be involved in cancer development. [provided by RefSeq, Jul 2008],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of various cells using Netrin-1 Monoclonal Antibody