



SR-6 Monoclonal Antibody

Catalog No	BYmab-12822
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	HTR6
Protein Name	5-hydroxytryptamine receptor 6
Immunogen	The antiserum was produced against synthesized peptide derived from human 5-HT-6. AA range:221-270
Specificity	SR-6 Monoclonal Antibody detects endogenous levels of SR-6 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
	T mg/m
Purity	≥90%
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Purity Storage Stability Synonyms	≥90% -20°C/1 year HTR6; 5-hydroxytryptamine receptor 6; 5-HT-6; 5-HT6; Serotonin receptor 6
Purity Storage Stability Synonyms Observed Band	 ≥90% -20°C/1 year HTR6; 5-hydroxytryptamine receptor 6; 5-HT-6; 5-HT6; Serotonin receptor 6 47kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	 ≥90% -20°C/1 year HTR6; 5-hydroxytryptamine receptor 6; 5-HT-6; 5-HT6; Serotonin receptor 6 47kD Cell membrane; Multi-pass membrane protein. Expressed in several human brain regions, most prominently in the caudate
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	 ≥90% -20°C/1 year HTR6; 5-hydroxytryptamine receptor 6; 5-HT-6; 5-HT6; Serotonin receptor 6 47kD Cell membrane; Multi-pass membrane protein. Expressed in several human brain regions, most prominently in the caudate nucleus. function:This is one of the several different receptors for 5-hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. The activity of this receptor is mediated by G proteins that stimulate adenylate cyclase. It has a high affinity for tricyclic psychotropic drugs.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in several human brain regions, most prominently in the




国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务,欢迎咨询

	AMP-dependent signaling pathway. This receptor is thought to regulate cholinergic neuronal transmission in the brain. Several antidepressants and antipsychotic drugs have a high affinity for this receptor. [provided by RefSeq, Aug 2013],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd