




Neurotensin Monoclonal Antibody

| Catalog No | BYmab-12864 |
|--------------------|---|
| Isotype | IgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | NTS |
| Protein Name | Neurotensin/neuromedin N |
| Immunogen | The antiserum was produced against synthesized peptide derived from the C-terminal region of human NTS. AA range:121-170 |
| Specificity | Neurotensin Monoclonal Antibody detects endogenous levels of Neurotensin protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | NTS; Neurotensin/neuromedin N |
| Observed Band | 20kD |
| Cell Pathway | Secreted. Cytoplasmic vesicle, secretory vesicle. Packaged within secretory vesicles. |
| Tissue Specificity | Brain,Tongue, |
| Function | function:Neurotensin may play an endocrine or paracrine role in the regulation of fat metabolism. It causes contraction of smooth muscle.,online information:Neurotensin entry,similarity:Belongs to the neurotensin family.,subcellular location:Packaged within secretory vesicles., |
| Background | This gene encodes a common precursor for two peptides, neuromedin N and neurotensin. Neurotensin is a secreted tridecapeptide, which is widely distributed throughout the central nervous system, and may function as a neurotransmitter or a neuromodulator. It may be involved in dopamine-associated pathophysiological events, in the maintenance of gut structure and function, and in the regulation of fat metabolism. Neurotensin also exhibits antimicrobial activity against bacteria |
| | |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








| | and fungi. Tissue-specific processing may lead to the formation in some tissues of larger forms of neuromedin N and neurotensin. The large forms may represent more stable peptides that are also biologically active. [provided by RefSeq, Oct 2014], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images



Western Blot analysis of various cells using Neurotensin Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd