



Stra8 Monoclonal Antibody

Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in		
Reactivity Applications WB Gene Name STRA8 Protein Name Immunogen	Catalog No	BYmab-12865
Applications Gene Name STRA8 Protein Name Stimulated by retinoic acid gene 8 protein homolog Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human STRA8. AA range:151-200 Specificity Stra8 Monoclonal Antibody detects endogenous levels of Stra8 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent . Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function: Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase. PTM:Phosphorylated ., tissue specificity:Expressed specifically in testis on the regulation of meiotic initiation in both spermadogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermadogenesis in	Isotype	IgG
Gene Name STRA8 Protein Name Stimulated by retinoic acid gene 8 protein homolog Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human STRA8. AA range:151-200 Specificity Stra8 Monoclonal Antibody detects endogenous levels of Stra8 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent . Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function: Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase., PTM:Phosphorylated, tissue specificity(Expressed specifically in testis.) Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in bospermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Reactivity	Human;Rat;Mouse;
Protein Name Stimulated by retinoic acid gene 8 protein homolog	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from the Internal region of human STRA8. AA range:151-200 Specificity Stra8 Monoclonal Antibody detects endogenous levels of Stra8 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophasePTM:Phosphorylated.,tissue specificity:Expressed specifically in testis. Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in espermatogenesis in	Gene Name	STRA8
Internal region of human STRA8. AA range:151-200 Specificity Stra8 Monoclonal Antibody detects endogenous levels of Stra8 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent. Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase, PTM:Phosphorylated., tissue specificity:Expressed specifically in testis. Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought fhat this gene may play a role in spermatogenesis in	Protein Name	Stimulated by retinoic acid gene 8 protein homolog
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent. Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function: Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase. PTM: Phosphorylated., tissue specificity: Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Specificity	Stra8 Monoclonal Antibody detects endogenous levels of Stra8 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent . Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase. PTM:Phosphorylated., tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent . Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Concentration	1 mg/ml
Synonyms STRA8; Stimulated by retinoic acid gene 8 protein homolog Observed Band 66kD Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Storage Stability	-20°C/1 year
Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Synonyms	STRA8; Stimulated by retinoic acid gene 8 protein homolog
Tissue Specificity Expressed specifically in testis and fetal ovaries. Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Observed Band	66kD
Function function:Required for the transition into meiosis for both female and male germ cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Cell Pathway	Cytoplasm . Nucleus . Shuttles between nucleus and cytoplasm. Nuclear export is XPO1-dependent
cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue specificity:Expressed specifically in testis., Background This gene encodes a retinoic acid-responsive protein. A homologous protein in mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Tissue Specificity	Expressed specifically in testis and fetal ovaries.
mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in	Function	cells. In female germ cells, required for premeiotic DNA replication and subsequent events in meiotic prophase.,PTM:Phosphorylated.,tissue
	Background	mouse has been shown to be involved in the regulation of meiotic initiation in both spermatogenesis and oogenesis, though feature differences between the mouse and human proteins suggest that these homologs are not entirely functionally equivalent. It is thought that this gene may play a role in spermatogenesis in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Stra8 Monoclonal Antibody