



NTN Monoclonal Antibody

| Catalog No | BYmab-12866 |
|--------------------|--|
| lsotype | lgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | NRTN |
| Protein Name | Neurturin |
| Immunogen | The antiserum was produced against synthesized peptide derived from the Internal region of human NRTN. AA range:111-160 |
| Specificity | NTN Monoclonal Antibody detects endogenous levels of NTN protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | NRTN; Neurturin |
| Observed Band | 22kD |
| Cell Pathway | Secreted. |
| Tissue Specificity | Melanoma,Peripheral blood lymphocyte, |
| Function | disease:Defects in NRTN are a cause of Hirschsprung disease (HSCR) [MIM:142623]. In association with mutations of RET gene, and possibly with other loci, defects in NRTN are involved in Hirschsprung disease. This genetic disorder of neural crest development is characterized by the absence of intramural ganglion cells in the hindgut, often resulting in intestinal obstruction.,function:Supports the survival of sympathetic neurons in culture. May regulate the development and maintenance of the CNS. Might control the size of non-neuronal cell population such as haemopoietic cells.,similarity:Belongs to the TGF-beta family. GDNF subfamily.,subunit:Homodimer; disulfide-linked., |

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| neurturin(NRTN) Homo sapiens This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. The encoded preproprotein is proteolytically processed to generate the mature protein. This protein signals through the RET receptor tyrosine kinase and a GPI-linked coreceptor, and promotes survival of neuronal populations. A neurturin mutation has been described in a family with Hirschsprung Disease. [provided by RefSeq, Aug 2016], |
|---|
| Avoid repeated freezing and thawing! |
| This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| |

Products Images



Western Blot analysis of various cells using NTN Monoclonal Antibody

Nanjing BYabscience technology Co.,Ltd