



# NOTCH4 (Cleaved-Val1467) mouse mAb

<b>Catalog No</b>	BYmab-12893
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Monkey
<b>Applications</b>	WB
<b>Gene Name</b>	NOTCH4 INT3
<b>Protein Name</b>	NOTCH4 (Cleaved-Val1467)
<b>Immunogen</b>	Synthesized peptide derived from human NOTCH4 (Cleaved-Val1467)
<b>Specificity</b>	This antibody detects endogenous levels of Human,Monkey NOTCH4 (Cleaved-Val1467, protein was cleaved amino acid sequence between 1466-1467 )
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal, Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Neurogenic locus notch homolog protein 4 (Notch 4;hNotch4) [Cleaved into: Notch 4 extracellular truncation; Notch 4 intracellular domain]
<b>Observed Band</b>	60 220kD
<b>Cell Pathway</b>	Cell membrane; Single-pass type I membrane protein.; [Notch 4 intracellular domain]: Nucleus. Following proteolytical processing NICD is translocated to the nucleus.
<b>Tissue Specificity</b>	Highly expressed in the heart, moderately in the lung and placenta and at low levels in the liver, skeletal muscle, kidney, pancreas, spleen, lymph node, thymus, bone marrow and fetal liver. No expression was seen in adult brain or peripheral blood leukocytes.
<b>Function</b>	alternative products:Experimental confirmation may be lacking for some isoforms,function:Functions as a receptor for membrane-bound ligands Jagged1, Jagged2 and Delta1 to regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBP-J kappa and activates genes of the enhancer of split locus. Affects the implementation of differentiation, proliferation and apoptotic

Nanjing BYabscience technology Co.,Ltd



programs. May regulate branching morphogenesis in the developing vascular system., polymorphism: The poly-Leu region of NOTCH4 (in the signal peptide) is polymorphic and the number of Leu varies in the population (from 6 to 12)., PTM: Phosphorylated., PTM: Synthesized in the endoplasmic reticulum as an inactive form which is proteolytically cleaved by a furin-like convertase in the trans-Golgi network before it reaches the plasma

#### Background

notch 4 (NOTCH4) Homo sapiens This gene encodes a member of the NOTCH family of proteins. Members of this Type I transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple different domain types. Notch signaling is an evolutionarily conserved intercellular signaling pathway that regulates interactions between physically adjacent cells through binding of Notch family receptors to their cognate ligands. The encoded preprotein is proteolytically processed in the trans-Golgi network to generate two polypeptide chains that heterodimerize to form the mature cell-surface receptor. This receptor may play a role in vascular, renal and hepatic development. Mutations in this gene may be associated with schizophrenia. Alternative splicing results in multiple transcript variants, at least one of which

#### matters needing attention

Avoid repeated freezing and thawing!

#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images