



CysLTR2 Monoclonal Antibody

Catalog No	BYmab-13192
lsotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	CYSLTR2
Protein Name	Cysteinyl leukotriene receptor 2
Immunogen	The antiserum was produced against synthesized peptide derived from human CYSLTR2. AA range:281-330
Specificity	CysLTR2 Monoclonal Antibody detects endogenous levels of CysLTR2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
,	
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year CYSLTR2; CYSLT2; CYSLT2R; PSEC0146; Cysteinyl leukotriene receptor 2; CysLTR2; G-protein coupled receptor GPCR21; hGPCR21; G-protein coupled receptor HG57; HPN321
Storage Stability Synonyms Observed Band	-20°C/1 year CYSLTR2; CYSLT2; CYSLT2R; PSEC0146; Cysteinyl leukotriene receptor 2; CysLTR2; G-protein coupled receptor GPCR21; hGPCR21; G-protein coupled receptor HG57; HPN321 40kD
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year CYSLTR2; CYSLT2; CYSLT2R; PSEC0146; Cysteinyl leukotriene receptor 2; CysLTR2; G-protein coupled receptor GPCR21; hGPCR21; G-protein coupled receptor HG57; HPN321 40kD Cell membrane; Multi-pass membrane protein.
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year CYSLTR2; CYSLT2; CYSLT2R; PSEC0146; Cysteinyl leukotriene receptor 2; CysLTR2; G-protein coupled receptor GPCR21; hGPCR21; G-protein coupled receptor HG57; HPN321 40kD Cell membrane; Multi-pass membrane protein. Widely expressed, with highest levels in the heart, placenta, spleen, peripheral blood leukocytes and adrenal gland. In lung, expressed in the interstitial macrophages, and slightly in smooth muscle cells.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Background	The cysteinyl leukotrienes LTC4, LTD4, and LTE4 are important mediators of human bronchial asthma. Pharmacologic studies have determined that cysteinyl leukotrienes activate at least 2 receptors, the protein encoded by this gene and CYSLTR1. This encoded receptor is a member of the superfamily of G protein-coupled receptors. It seems to play a major role in endocrine and cardiovascular systems. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

expressed in the interstitial macrophages, and slightly in smooth muscle cells.,

Products Images



Nanjing BYabscience technology Co.,Ltd