





Flg Monoclonal Antibody

Catalog No BYmab-13244 Isotype IgG Reactivity Human;Mouse;Rat Applications WB Gene Name FGFR1 Protein Name Fibroblast growth factor receptor 1 Immunogen The antiserum was produced against synthesized peptide derived from human FGFR1. AA range:628-675 Specificity FIg Monoclonal Antibody detects endogenous levels of Flg protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein, Nucleus, Cytoplasmic versicle. After ligand binding, both receptor and ligand rapidly internalized. Can translocate to		
Reactivity Human;Mouse;Rat Applications WB Gene Name FGFR1 Protein Name Fibroblast growth factor receptor 1 Immunogen The antiserum was produced against synthesized peptide derived from human FGFR1. AA range:626-675 Specificity Fig Monoclonal Antibody detects endogenous levels of Flg protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus in foreskin fibr	Catalog No	BYmab-13244
Applications WB Gene Name FGFR1 Protein Name Fibroblast growth factor receptor 1 Immunogen The antiserum was produced against synthesized peptide derived from human FGFR1. AA range:626-675 Specificity Fig Monoclonal Antibody detects endogenous levels of Fig protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Isotype	IgG
Gene Name FGFR1 Protein Name Fibroblast growth factor receptor 1 Immunogen The antiserum was produced against synthesized peptide derived from human FGFR1. AA range:626-675 Specificity Flg Monoclonal Antibody detects endogenous levels of Flg protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma an	Reactivity	Human;Mouse;Rat
Protein Name Fibroblast growth factor receptor 1 Immunogen The antiserum was produced against synthesized peptide derived from human FGFR1. AA range:626-675 Specificity FIg Monoclonal Antibody detects endogenous levels of FIg protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fi	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human FGFR1. AA range:626-675 Specificity Fig Monoclonal Antibody detects endogenous levels of Flg protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Gene Name	FGFR1
FGFR1. AA range:626-675 Specificity FIg Monoclonal Antibody detects endogenous levels of FIg protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adreal cortex cell lines. Some isoform 17 are not detected in these cells.	Protein Name	Fibroblast growth factor receptor 1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type 1 membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Specificity	Flg Monoclonal Antibody detects endogenous levels of Flg protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type 1 membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Concentration	1 mg/ml
FGFR1; BFGFR; CEK; FGFBR; FLG; FLT2; HBGFR; Fibroblast growth factor receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Purity	≥90%
receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen CD331 Observed Band 91868DA Cell Pathway Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Storage Stability	-20°C/1 year
Cell membrane; Single-pass type I membrane protein. Nucleus. Cytoplasm, cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Synonyms	receptor 1; FGFR-1; Basic fibroblast growth factor receptor 1; BFGFR; bFGF-R-1; Fms-like tyrosine kinase 2; FLT-2; N-sam; Proto-oncogene c-Fgr; CD antigen
cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol, and from there to the nucleus. Tissue Specificity Detected in astrocytoma, neuroblastoma and adrenal cortex cell lines. Some isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Observed Band	91868DA
isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform 18 and isoform 19 are not detected in these cells.	Cell Pathway	cytosol. Cytoplasmic vesicle. After ligand binding, both receptor and ligand are rapidly internalized. Can translocate to the nucleus after internalization, or by translocation from the endoplasmic reticulum or Golgi apparatus to the cytosol,
Function	Tissue Specificity	isoforms are detected in foreskin fibroblast cell lines, however isoform 17, isoform
	Function	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images