





FPR1 Monoclonal Antibody

Catalog No BYmab-13250 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name FPR1 Protein Name fMet-Leu-Phe receptor Immunogen The antiserum was produced against synthesized peptide derived from human FPR1. AA range:151-200 Specificity FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function-High affinity receptor for N-formy		
Reactivity Applications WB Gene Name FPR1 Protein Name Immunogen The antiserum was produced against synthesized peptide derived from human FPR1. AA range:151-200 Specificity FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dillution WB 1:500-2000 Concentration 1 mg/ml 290% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a 6-protein that activates a phosphatidylinositol-calcium second messenger system. PTM:Phosphorylated; which is necessary for desensitization, similarity:Belongs to the G-protein coupled receptor 1 family, tissue specificity:Neutrophils. This gene encodes a G protein-coupled receptor of mammalian phagocyctic cells that is a member of the	Catalog No	BYmab-13250
Applications WB Gene Name FPR1 Protein Name fMet-Leu-Phe receptor Immunogen The antiserum was produced against synthesized peptide derived from human FPR1. AA range:151-200 Specificity FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemodactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a 6-prosein-toughed in the receptor causes along the prosein-coupled receptor of member of the receptor from the dictivates a phosphaticlylinositol-calcium second messenger system. PTM:Phosphorylated; which is necessary for desensitizating, similarity Belongs to the G-protein coupled receptor of family, tissue specificity. Neutrophils. Background for the protein coupled receptor of mammalian phagocycic cells that is a member of the	Isotype	IgG
Gene Name FPR1 Protein Name fMet-Leu-Phe receptor Immunogen The antiserum was produced against synthesized peptide derived from human FPR1. AA range:151-200 Specificity FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane : Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemoatctic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. PTM:Phosphorylated; which is necessary for desensitization. similarity:Belongs to the G-protein coupled receptor 1 (FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Reactivity	Human;Rat;Mouse;
Protein Name fMet-Leu-Phe receptor Immunogen The antiserum was produced against synthesized peptide derived from human FPR1. AA range:151-200 Specificity FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, lgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor of mammalian phagocytic cells that is a member of the Background formyl peptide receptor of fPPR1) Homo sapiens This gene encodes a G protein the <th>Applications</th> <th>WB</th>	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human FPR1. AA range:151-200 Specificity FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG. Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization. Background	Gene Name	FPR1
FPR1. AA range:151-200 Specificity FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG. Tissue Specificity Neutrophils. Function function: High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinosiolo-calcium second messenger system. PfM:Phosphorylated; which is necessary for desensitization, similarity. Belongs to the G-protein coupled receptor 1 family. tissue specificity. Neutrophiles. Background formyl peptide receptor of mammalian phagocytic cells that is a member of the	Protein Name	fMet-Leu-Phe receptor
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG. Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. PTM:Phosphorylated; which is necessary for desensitization., similarity:Belongs to the G-protein coupled receptor 1 (FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Immunogen	
Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1(FPR1) Homo sapiens This gene encodes a G protein-coupled receptor for mammalian phagocytic cells that is a member of the	Specificity	FPR1 Monoclonal Antibody detects endogenous levels of FPR1 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane ; Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function: High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. PTM: Phosphorylated; which is necessary for desensitization., similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Neutrophils. Background formyl peptide receptor of mammalian phagocytic cells that is a member of the	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG. Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family,tissue specificity:Neutrophils. Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G-protein-coupled receptor of mammalian phagocytic cells that is a member of the	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils. Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG. Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Dilution	WB 1:500-2000
Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell Pathway Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function: High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G-protein-coupled receptor of mammalian phagocytic cells that is a member of the	Concentration	1 mg/ml
Synonyms FPR1; fMet-Leu-Phe receptor; fMLP receptor; N-formyl peptide receptor; FPR; N-formylpeptide chemoattractant receptor Observed Band 38kD Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function: High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Purity	≥90%
N-formylpeptide chemoattractant receptor 38kD Cell Pathway Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G-protein-coupled receptor of mammalian phagocytic cells that is a member of the	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG Tissue Specificity Neutrophils. Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G-protein-coupled receptor of mammalian phagocytic cells that is a member of the	Synonyms	
ligands, fMLP, TAFA4 and CTSG Neutrophils. Function function: High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Observed Band	38kD
Function function:High affinity receptor for N-formyl-methionyl peptides, which are powerful neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Cell Pathway	Cell membrane ; Multi-pass membrane protein . Internalizes in presence of its ligands, fMLP, TAFA4 and CTSG
neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Neutrophils., Background formyl peptide receptor 1(FPR1) Homo sapiens This gene encodes a G protein-coupled receptor of mammalian phagocytic cells that is a member of the	Tissue Specificity	Neutrophils.
protein-coupled receptor of mammalian phagocytic cells that is a member of the	Function	neutrophils chemotactic factors. Binding of FMLP to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system.,PTM:Phosphorylated; which is necessary for desensitization.,similarity:Belongs to the G-protein coupled
	Background	protein-coupled receptor of mammalian phagocytic cells that is a member of the

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	phagocytic cells to invasion of the host by microorganisms and is important in host defense and inflammation.[provided by RefSeq, Jul 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658