



Glucagon Receptor Monoclonal Antibody

| Catalog No | BYmab-13275 |
|--------------------|--|
| Isotype | lgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB |
| Gene Name | GCGR |
| Protein Name | Glucagon receptor |
| Immunogen | The antiserum was produced against synthesized peptide derived from human GLR. AA range:95-144 |
| Specificity | Glucagon Receptor Monoclonal Antibody detects endogenous levels of Glucagon Receptor protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Monoclonal, Mouse,IgG |
| Purification | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | GCGR; Glucagon receptor; GL-R |
| Observed Band | 54kD |
| Cell Pathway | Cell membrane ; Multi-pass membrane protein . Is rapidly internalized after ligand-binding |
| Tissue Specificity | Liver,Placenta, |
| Function | disease: It is a candidate for certain defects in non-insulin-dependent diabetes mellitus (NIDDM). The variant in position 40 (Ser) is found in some NIDDM patients, but also in nondiabetic subjects.,function: This is a receptor for glucagon which plays a central role in regulating the level of blood glucose by controlling the rate of hepatic glucose production and insulin secretion. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase and also a phosphatidylinositol-calcium second messenger system.,similarity:Belongs to the G-protein coupled receptor 2 family., |
| Background | glucagon receptor(GCGR) Homo sapiens The protein encoded by this gene is a glucagon receptor that is important in controlling blood glucose levels. Defects in |
| | Nanjing BYabscience technology Co.,Ltd |

Brabscience technology Co.,Ltd Nanjing

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| | this gene are a cause of non-insulin-dependent diabetes mellitus (NIDDM).[provided by RefSeq, Jan 2010], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |

Products Images



Nanjing BYabscience technology Co.,Ltd