



GPR87 Monoclonal Antibody

Catalog No BYmab-13352 Isotype IgG Reactivity Human;Mouse Applications WB Gene Name GPR87 Protein Name G-protein coupled receptor 87 Immunogen The antiserum was produced against synthesized peptide derived from human GPR87. AA range:221-270 Specificity GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in trainily, tissue specificity. Expressed on placenta and prostate. We		
Reactivity Human;Mouse Applications WB Gene Name GPR87 Protein Name G-protein coupled receptor 87 Immunogen The antiserum was produced against synthesized peptide derived from human GPR87. AA range;221-270 Specificity GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein . Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in squamous cell carcinoma of the lung. Function function, Orphan receptor, similarity, Belongs to the G-protein coupled receptor 1 family, tissue specificity/Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in througaumonous cell carcinoma of PMID:18057535)	Catalog No	BYmab-13352
Applications WB Gene Name GPR87 Protein Name G-protein coupled receptor 87 Immunogen The antiserum was produced against synthesized peptide derived from human GPR87. AA range:221-270 Specificity GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in squamous cell carcinoma of the lung. Function Unction, Orphan receptor, similarity. Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in ung squamous cell carcinoma (PMID: 18057635)	Isotype	IgG
Gene Name GPR87 Protein Name G-protein coupled receptor 87 Immunogen The antiserum was produced against synthesized peptide derived from human GPR87. AA range:221-270 Specificity GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in a dumity, tissue specificity: Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Function This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encode	Reactivity	Human;Mouse
Protein Name G-protein coupled receptor 87 Immunogen The antiserum was produced against synthesized peptide derived from human GPR87. AA range:221-270 Specificity GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor., similarity:Belongs to the G-protein coupled receptor 1 family, tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Background This gene encodes a G pro	Applications	WB
Immunogen The antiserum was produced against synthesized peptide derived from human GPR87. AA range:221-270 Specificity GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family. dissue specificity: Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromos	Gene Name	GPR87
Specificity GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor, similarity:Belongs to the G-protein coupled receptor 1 family, tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Protein Name	G-protein coupled receptor 87
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Monoclonal, Mouse,IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function: Orphan receptor. similarity:Belongs to the G-protein coupled receptor 1 family. tissue specificity; Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Immunogen	
Source Monoclonal, Mouse, IgG Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID::18057535)	Specificity	GPR87 Monoclonal Antibody detects endogenous levels of GPR87 protein.
Purification The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family, tissue specificity: Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor., similarity:Belongs to the G-protein coupled receptor 1 family., tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMIID:18057535)	Source	Monoclonal, Mouse,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057635)	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function: Orphan receptor, similarity: Belongs to the G-protein coupled receptor 1 family., tissue specificity: Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Dilution	WB 1:500-2000
Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor., similarity:Belongs to the G-protein coupled receptor 1 family., tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Concentration	1 mg/ml
Synonyms GPR87; GPR95; FKSG78; G-protein coupled receptor 87; G-protein coupled receptor 95 Observed Band 40kD Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function: Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Purity	≥90%
Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Synonyms	
Tissue Specificity Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Observed Band	40kD
expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in squamous cell carcinoma of the lung. Function function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Cell Pathway	Cell membrane ; Multi-pass membrane protein .
family.,tissue specificity:Expressed in placenta and prostate. Weaker expression in thymus. Not expressed in thalamus, hippocampus, pons or cerebellum., Background This gene encodes a G protein-coupled receptor and is located in a cluster of G protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Tissue Specificity	expressed in thalamus, hippocampus, pons or cerebellum. Overexpressed in
protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)	Function	family, tissue specificity:Expressed in placenta and prostate. Weaker expression
	Background	protein-couple receptor genes on chromosome 3. The encoded protein has been shown to be overexpressed in lung squamous cell carcinoma (PMID:18057535)

Nanjing BYabscience technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

