



Histamine H2 Receptor Monoclonal Antibody

Catalog No	BYmab-13360
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	HRH2
Protein Name	Histamine H2 receptor
Immunogen	The antiserum was produced against synthesized peptide derived from human HRH2. AA range:131-180
Specificity	Histamine H2 Receptor Monoclonal Antibody detects endogenous levels of Histamine H2 Receptor protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HRH2; Histamine H2 receptor; H2R; HH2R; Gastric receptor I
Observed Band	40kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Brain,Liver,Skin,Stomach,
Function	function:The H2 subclass of histamine receptors mediates gastric acid secretion. Also appears to regulate gastrointestinal motility and intestinal secretion. Possible role in regulating cell growth and differentiation. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase and, through a separate G protein-dependent mechanism, the phosphoinositide/protein kinase (PKC) signaling pathway.,miscellaneous:Antagonists for this receptor have proven to be effective therapy for acid peptic disorders of the gastrointestinal tract. Certain antagonists are used in the treatment of neuropsychiatric and neurological diseases such as schizophrenia, Alzheimer disease and Parkinson disease.,online information:H2 receptor entry,similarity:Belongs to the G-protein coupled receptor 1 family.,

Nanjing BYabscience technology Co.,Ltd



Background

Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4. Histamine receptor H2 belongs to the family 1 of G protein-coupled receptors. It is an integral membrane protein and stimulates gastric acid secretion. It also regulates gastrointestinal motility and intestinal secretion and is thought to be involved in regulating cell growth and differentiation. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using Histamine H2 Receptor Monoclonal Antibody